

GENETIC ANALYSIS AND VALIDATION OF TRAIT SPECIFIC SSR MARKERS FOR YIELD AND ITS ATTRIBUTING TRAITS IN F₃ FAMILY OF IM 90/ IR 64 CROSS IN RICE (*ORYZA SATIVA* L.) UNDER AEROBIC CONDITION

POTLANNAGARI ROOPA SOWJANYA, P. RAGHAVENDRA * AND SHAILAJA HITTALMANI

Marker assisted selection laboratory,

Dept. of Genetics and plant breeding, UAS, Bangalore - 560 065, Karnataka, INDIA

e-mail: raghavendra9078@gmail.com

INTRODUCTION

Rice (*Oryza sativa* L.) is a plant belonging to the family of grasses, Gramineae. It is one of the three major food crops of the world and forms the staple diet of about half of the world's population. The global production of rice has been estimated to be at the level of 650 million tones. Asia is the leader in rice production accounting for about 90% of the world's production (FAOSTAT, 2013). Hybridization is one of the way through which enormous variability can be created (Shailaja Hittalmani and Shivashankar, 1987). It has been postulated that advancement in breeding programme depends on extent of variability created during hybridization (Shantavva *et al.*, 2014). Segregating populations are more important for improving plant types by operating selection (Farhad Kahani and Shailaja Hittalmani 2015). Therefore the acquaintance of genetic parameters present in segregating population for the improvement of character under study has paramount importance. Phenotypic coefficient of variation(PCV), genotypic co-efficient of variation (GCV), heritability, genetic advancement will be useful for efficient exploitation of variability (Shantavva *et al.*, 2014).The genotypic coefficient of variation measures the range of variability available in a crop and also enables to compare the amount of variability present in different characters. The phenotypic expression of the character is the result of interaction between genotype and environment. Hence, the total variance needs to be partitioned into heritable and non-heritable components to assess the inheritance pattern of the particular character under study. Heritability indicates the relative degree at which a character is transmitted from parent to offspring. Heritability estimates along with GA are normally more helpful in predicting the gain under selection than heritability estimates alone. As both will serve as important selection parameters (Ogunbayo *et al.*, 2014). One of the main constraints of rice cultivation and production is water shortage during periods of low rainfall, which affects the vegetative growth rate and grain yield (Manjappa *et al.*, 2014).Aerobic rice cultivation is one of the alternative wherein the water required is half of that of irrigated puddled rice, without compromising the productivity (Gandhi *et al.*, 2011a;Gandhi *et al.*, 2011b;Nareshbabu *et al.*, 2010).It has been reported that, validation of markers helps in determining the reliability and practical applicability of the markers in predicting the phenotype (Collard *et al.*, 2005).Among the markers SSR markers are considered as marker of choice,because it has the advantage of being co-dominant; they are simple PCR based and extremely polymorphic. They are also highly informative due to number and frequency of alleles detected and ability to distinguish between closely related individuals. They find application as marker for mapping, cultivar identification, protecting germplasm, determination of hybridity, analysis of gene pool variation and as diagnostic marker for traits of economic value (Powell *et al.*, 1996).In this context the present investigation was undertaken to evaluate the genetic variability present in F₃ segregating populations and also to validate trait specific SSR markers linked to yield and related traits in rice.

ABSTRACT

Thirty F₃ families derived from IM 90/ IR 64 cross in rice were evaluated under aerobic condition for 14 quantitative traits to assess the genetic parameters and validate SSR markers linked to yield related traits. The estimates of GCV were lower than the estimates of respective PCV, indicating the influence of environmental factors on the expression of the traits studied. Further the traits like spikelet fertility (94.99%, 14.62), plant height at maturity (91.22 %, 12.09) and Days to 50% flowering (81.45%,12.20) have high heritability coupled with moderate genetic advance as percent mean. Indicates greater role of additive gene action in their inheritance. The results of correlation revealed positive and significant correlation of grain yield per plant with productive tillers per plant (r=0.490), spikelet per panicle(r=0.422), harvest index (r=0.369) and biomass(r=0.294). Whereas, the highest direct effects on grain yield per plant by productive tillers per plant (1.083), biomass(2.76) and panicle weight(2.22) was observed indicated that during selection these traits has to be given prime importance. Further the six validated SSR markers associated with yield related traits could be useful for selection in segregating populations of rice.

KEY WORDS

Aerobic rice
F₃ family
Genetic parameters
SSR markers

Received : 07.01.2016

Revised : 22.02.2016

Accepted : 15.04.2016

*Corresponding author

MATERIALS AND METHODS

Plant material

The thirty F₃ families along with parents and checks were evaluated during *kharif* 2013, in augmented design (Federer 1965), under aerobic conditions in K-block, Dept. of Genetic and Plant Breeding, UAS, Bangalore which represents the eastern dry zone, with all proper crop management practices. The spacing was kept at row-row 20 cm and plant to plant 25 cm.

Aerobic rice cultivation

Aerobic rice cultivation involves direct sowing of seeds in unpuddled land with a spacing of 30 cm × 25 cm. Irrigation was given at 5 days interval to maintain soil moisture at field capacity throughout the crop growth. The soil is maintained under aerobic condition unlike the anaerobic condition in puddled soils with standing water in irrigated transplanted rice. During reproductive stage the irrigation is given once in 3 days interval.(Raghavendra and Shailaja Hittalmani 2015).

Observations recorded

In order to obtain the information on the performance of families, 20 randomly selected plant data was recorded for 14 quantitative traits as per SES 1996. Among those growth traits like plant height at 30days after transplanting, plant height at maturity, days to 50 % flowering, total number of tillers per plant and spikelets per panicle. Yield traits like productive tillers per plant, spikelet fertility, panicle length, panicle weight, grain yield per plant, 1000 grain weight, biomass, harvest index were recorded.

DNA extraction and SSR marker analysis

Leaf samples of twenty five days old seedlings were used for DNA extraction. DNA was isolated as Dellaporta method (Dellaporta, *et al.*, 1983).Total of 500 rice microsatellite markers used for screening IM 90 and IR 64 for polymorphism survey. Further, twenty reported SSR markers associated with yield and related traits, with high LOD score markers were used for genotyping of F₃ progenies (Table 1). PCR amplification was carried out following the standard procedures. The bands generated by polymorphic SSR primers were scored by giving code 'A' for Parent-1 (IM 90) (upper/ lower band), 'B' for Parent-2 (IR64) (upper/ lower band) and – for missing samples (Fig.1).

Statistical analysis

The data recorded on 20 randomly selected plants were subjected to ANOVA (Panse and Sukhatme, 1967). Variance, heritability and genetic advance (Robinson and Comstock, 1949) were estimated. Single marker analysis was done with the help of "student's" t-distribution and one-way analysis of variance (Edwards *et al.*, 1987).

RESULTS AND DISCUSSION

ANOVA

The mean sum of squares due to families showed significant difference for 14 quantitative traits (Table 2) indicates the presence of adequate genetic variability (Paikhomba *et al.*, 2014). This would prove to be beneficial as available variability could be further utilized in crop improvement programme.

Table 1: List of SSR trait specific markers used for validation for yield and its contributing traits in Rice

Sl. No	Marker	SSR marker linked traits	Chromo some No.	Forward primer 5' - 3'	Reverse primer 5' - 3'	Expected base pair	References
1	RM24	Plant height (cm)	1	GAAGTGTGATCACTGTAACC	TACAGTGGACGGCGAAGTCC	193	Price <i>et al.</i> (2002)
2	RM 243	Root dry weight (g)	1	GATCTGCAGACTGCAGTTGC	AGCTGCAACGATGTTGTCC	116	Marriet <i>et al.</i> (2005)
3	RM443	Test weight (g)	1	GATGGTTTCATCGGCTACC	AGTCCAGAAATGTCGTTCC	124	Price <i>et al.</i> (2002)
4	RM493	Spikelet fertility (%)	1	TAGTCCAAACAGGATCGACC	GTACGTAACCGGGAAAGGTG	211	Lauxmanet <i>al.</i> (2010)
5	RM495	Plant height (cm), spikelet fertility (%)	1	AATCCAAGGTGCAGAGATGG	CAACGATGACGAACAACACC	159	Yueet <i>al.</i> (2005)
6	RM594	Root to shoot ratio	1	GCCACCAGTAAAGCAATAC	TTGATCTGCTAGTGAGACCC	300	Marriet <i>al.</i> (2005)
7	RM208	Panicle length (cm), number of spikelets per panicle	2	TCTGCAAGCCTTGCTGATG	TAAGTCGATCATTTGTGGACC	173	Venuprasadet <i>al.</i> (2009)
8	RM263	Grain yield (stress) (g)	2	CCCAGGCTAGCTCATGAACC	GCTACGTTTGAGTACCACC	207	Zhang <i>et al.</i> (2001)
9	RM131	Grain yield (g)	4	TCCTCCCTCCCTTCGCCCACTG	CGATGTTCCGATGGCTGTCC	215	Yueet <i>al.</i> (2006)
10	RM471	Days to 50% flowering	4	ACGCACAAGCAGATGATGAG	GGGAGAAGACGAATGTTGC	106	Marriet <i>al.</i> (2005)
11	RM248	Root dry weight (g), Total root weight (g), days to 50% flowering	7	TCCTTGTGAAATCTGGTCCC	GTAGCCTAGCAATGGTGCATG	102	Price <i>et al.</i> (2002)
12	RM152	Spikelet fertility (%)	8	GAACCACCACACCTCACCG	CCGTAGACCTTCTTGAAGTAG	NA	Xing <i>et al.</i> (2002)
13	RM337	Plant height (cm)	8	GTAGGAAAGGAAAGGGCAGAG	CGATAGATAGTAGATGTGGCC	192	Lauxmanet <i>al.</i> (2010)
14	RM447	Test weight (g)	8	CCCTTGCTGTCTCCTCTC	ACGGCTTCTTCTCCTCTC	111	Lauxmanet <i>al.</i> (2010)
15	RM3262	Grain yield (g)	8	CTGGAGATGCAGATCTCAACC	TAGTACAACATGGGAGCCTGTCC	290	Laffite <i>et al.</i> (2002)
16	RM205	Total number of tillers	9	CTGGTCTGTATGGGAGCAG	CTGGCCCTCACGTTTCAGTG	222	Lauxmanet <i>al.</i> (2010)
17	RM242	Grain yield (g), root length (cm)	9	GGCAACGTTGTATGTCTC	TATATGCCAAGACGGATGGG	225	Marriet <i>al.</i> (2005)
18	RM596	Grain yield (stress) (g)	10	ATCTACACGGACGAATTGCC	AGAAGCTTCAGCCTCTGCAG	188	Lauxmanet <i>al.</i> (2010)
19	RM4	Productive tillers	12	TTGACGAGGTCAGCACTGAC	AGGGTGTATCCGACTCATCC	NA	Venuprasadet <i>al.</i> (2009)
20	RM519	Plant height (cm), productive tillers	12	AGAGAGCCCTAAATTTCCG	AGGTACCGTCACTGTGGAC	122	Venuprasadet <i>al.</i> (2009)

Table 2: Analysis of variance (mean sum of squares) for 14 quantitative characters studied in F₃ families of IM 90 / IR 64 Rice cross

Source of variation	Progenies	Error
Degrees of Freedom (df)	29	10
Days to 50% flowering	28.39***	3.63
Plant height at 30 days after transplanting (cm)	10.05***	5.18
Plant height at maturity (cm)	16.39***	0.98
Productive tillers plant ⁻¹	3.97*	1.58
Total number of tillers plant ⁻¹	1.59*	0.43
Spikelets per panicle	6.15***	5.15
Spikelet fertility (%)	3.17***	1.12
Panicle length (cm)	3.44*	1.21
Panicle weight (g)	0.53*	0.38
Days to maturity	50.96*	25.26
Grain yield plant ⁻¹ (g)	12.54*	5.16
1000-grain weight (g)	4.1*	0.64
Biomass (g)	53.54**	7.11
Harvest index	0.00*	0.00




Figure 1: Characterization of F₃ progenies of IM 90 X IR 64 cross in rice using SSR markers RM 248 linked to QTL's governing grain yield and RM 205 linked to QTL's governing number of tillers per plant.

Table 3: Estimates of genetic parameters for yield and its attributing traits in F₃ families of IM 90 / IR 64 of rice cross

Sl. No.	Plant traits	Range		Mean	PCV%	GCV%	h ² (bs)%	GAM
		Max.	Min.					
1	Days to 50 % flowering	116.00	94.65	103.37	4.28	3.86	81.45	12.20
2	Plant height at 30 DAT (cm)	40.80	29.70	35.42	17.50	16.55	76.25	15.11
3	Plant height at maturity (cm)	74.00	59.75	66.83	15.02	14.79	91.22	12.09
4	Productive tillers per plant	13.00	8.75	10.50	10.40	8.05	73.14	11.10
5	Total tillers per plant	15.00	8.00	12.15	8.89	7.07	63.24	14.84
6	Spikelets per panicle	94.30	85.15	90.23	2.24	2.14	81.32	15.40
7	Spikelet fertility (%)	88.58	80.88	84.66	1.84	1.79	94.99	14.62
8	Panicle length (cm)	21.00	15.95	18.56	15.83	13.77	41.90	6.45
9	Panicle weight (g)	5.00	1.06	3.26	21.33	19.65	20.48	11.53
10	Days to maturity	146.60	116.25	132.66	14.87	13.06	39.60	17.09
11	Grain yield per plant (g)	46.45	21.70	29.10	20.79	17.45	47.66	19.58
12	1000 grain weight (g)	29.92	20.84	22.25	17.63	16.72	51.08	17.56
13	Biomass (g)	82.75	37.60	63.63	21.50	19.33	32.75	21.45
14	Harvest index	0.52	0.31	0.47	16.22	11.59	44.32	18.87

Min. Minimum
 Max. Maximum
 SE Standard error
 h²(bs)% GAM
 Broad sense heritability
 Genetic advance as percent mean
 PCV (%) Phenotypic coefficient of variation (%)
 GCV (%) Genotypic coefficient of variation (%)

The estimates of genetic parameters like mean, coefficient of variation, heritability and genetic advance as percent mean deserve attention in deciding selection criteria for improvement in the concerned characters. (Debnath *et al.*, 2015)

Mean and Range

The mean performance of families for different traits are noticed as follows days to fifty per cent flowering ranged between 94.65 to 116 days with a mean of 103.37 days, Plant height at 30 days after transplanting varied between 29.70 to 40.80 cm with a mean value of 35.42 cm, plant height varied from 59.75 to 74.00 cm with a mean value of 66.83 cm, productive tillers per plant varied from 8.75 to 13.00 with a mean value of 10.50, total number of tillers per plant varied between 8.00 to 15.00 with a mean value of 12.15. Spikelets per panicle varied from 85.15 to 94.30 with a mean value of 90.23 cm, spikelet fertility ranged from 80.00 to 88.58% with a mean value of 84.66%, panicle length ranged from 15.95 to 21.00 cm with a mean value of 18.56cm, panicle weight varied from 1.06 to 5.00 g with a mean value of 3.26 g.

Days to maturity varied between 116.25 to 146.30 days after transplanting with a mean value of 132.66 days, grain yield per plant varied between 21.70 to 46.45 g with a mean value of 29.10 g, 1000 grain weight varied between 20.84 to 29.92 g with a mean value of 22.25 g, biomass ranged from 37.60 to 82.75 g with a mean value of 63.63 g, harvest index varied from 0.31 to 0.72 with a mean value of 0.47%. This finding corroborated with Kumar *et al.*, 2015

PCV and GCV

The estimates of GCV were lower than the PCV estimates (Khalid *et al.*, 2012 and Gangashetty *et al.*, 2013, Paikhombha *et al.*, 2014) indicating there is influence of environmental factors on the expression of traits (Table 3). The PCV estimates ranged from 1.84 to 21.50% and GCV estimates ranged from 1.79 to 19.65%.

Highest GCV was recorded for panicle weight (19.65) followed by Biomass (19.33), grain yield per plant (17.45) and 1000 grain weight (16.72), highest PCV is recorded for traits are Biomass (21.50), panicle weight (21.33), grain yield per plant (20.79), and 1000 grain weight (17.63) indicated presence of

Table 4: Correlation and direct effects of 14 quantitative traits on grain yield per plant of families derived from IM 90/IR 64 cross of rice

Traits	Correlation	Direct effects
Days to 50% flowering	-0.056	-0.1884
Plant height at 30 days after transplanting (cm)	0.450*	-0.0601
Plant height at maturity (cm)	0.358*	0.0530
Productive tillers plant ⁻¹	0.490*	1.0830
Total number of tillers plant ⁻¹	-0.129	0.0759
Spikelets per panicle	0.422*	0.0086
Spikelet fertility (%)	0.458*	0.0646
Panicle length (cm)	-0.026	-0.0431
Panicle weight (g)	0.012	2.2292
Days to maturity	0.200	0.1761
1000-grain weight (g)	0.553**	-0.0302
Biomass (g)	0.294*	2.7658
Harvest index	0.369*	-0.1422

predicting the resultant effect of selection on phenotypic expression, without genetic advance the estimates heritability will not be of practical use.

Heritability and genetic advance

The data on heritability (bs), depicted that heritability estimates % for 14 quantitative traits, under study ranged from 20.48% to 94.99%. Highest value of heritability was noticed for spikelet fertility (94.99%), followed by plant height at maturity (91.22%) and days to 50 % flowering (81.45%) indicated least influence of environment on these traits. Whereas, Least was observed for panicle weight (20.48), followed by biomass (32.75%) and harvest index (44.32%) indicated higher influence of environment on expression of these traits (Prajapati *et al.*, 2011).

Genetic advance on percent mean was high for biomass

Table 5: R² values for SSR markers showing significant association with different phenotypic characters in rice

Sl. No.	Marker	Chromosome No.	Trait linked to SSR marker	t value (2 tailed)	R ²
1.	RM495	1	Spikelet fertility (%)	2.576*	19.73%
			Plant height (cm)	2.172*	18.15%
2.	RM242	9	Grain yield (g)	1.493*	15.31%
3.	RM152	8	Spikelet fertility (%)	2.029*	12.06%
4.	RM263	2	Grain yield (g)	1.993*	12.04%
5	RM 248	7	Days to 50% flowering	1.162*	10.23%
6	RM596	10	Grain yield (g)	2.216*	4.45%

*significance at 5%, **significance at 1% and *** significance at 0.1% level

Table 6: Number of genotypes carrying positive alleles for the traits attributing to yield

Sl. No.	SSR Markers	Traits linked	Positive alleles carrying genotypes like IM 90 parent
1	RM4	Productive tillers	5
2	RM248	Root dry weight (g), Total root weight (g), days to 50% flowering	5
3	RM131	Grain yield (g)	7
4	RM596	Grain yield (stress) (g)	7
5	RM242	Grain yield (g), root length (cm)	8 and 4
6	RM263	Grain yield (stress) (g)	8
7	RM24	Plant height (cm)	10
8	RM152	Spikelet fertility (%)	10
9	RM443	Test weight (g)	10
10	RM495	Plant height (cm) and spikelet fertility (%)	10 and 7
11	RM519	Plant height (cm), productive tillers	10
12	RM205	Total number of tillers	11
13	RM493	Spikelet fertility (%)	11
14	RM447	Test weight (g)	14

high variability for these traits (Bidhan *et al.*, 2011; Ghosh and Sharma, 2012).

Least GCV was recorded for spikelet fertility (1.79), spikelets per panicle (2.14) and days to 50 % flowering (3.86). However, least PCV values are observed for spikelet fertility (1.84), spikelets per panicle (2.24) and days to 50 % flowering (4.28) indicated presence of low variability for these traits (Table.3). GCV along with heritability gave an idea of expected genetic gain from selection (Kumar *et al.*, 2009). Although GCV and PCV is the indicative of the presence of degree of genetic variation, which the amount of heritability portion of variation can only be determined (Bidhan *et al.*, 2011; Idris *et al.*, 2012). In

(21.45) followed by Grain yield per plant (19.58) and harvest index (18.87) indicated preponderance of additive gene action (Bidhan *et al.*, 2011). However, Least was observed for panicle length (6.45) followed by productive tillers per plant (11.10) and Panicle weight (11.53) indicated majority of non-additive gene action in these traits inheritance.

Most of the characters under study were having moderate-high heritability coupled with moderate-high genetic advance as percent of mean indicated the preponderance of additive gene effects in its inheritance and such characters could be improved by mass selection and other breeding methods based on progeny testing (Kumar *et al.*, 2015). However,

panicle length showed moderate values of heritability coupled with low genetic advance also indicated presence of additive gene action suggest that The high heritability is being exhibited due to favorable influence of environment rather than genotype and selection for such traits may not be rewarding. (Kumar *et al.*, 2015). Whereas for panicle weight low values of heritability coupled with moderate genetic advance was observed reveals that the trait is governed by additive gene effects. The low heritability is exhibited due to high environmental effects so selection may be effective in this case (Debnath *et al.*, 2015)

Correlation and path analysis

The results indicated positive and significant correlation of grain yield per plant with plant height at 30 days after transplanting ($r=0.450^*$), plant height at maturity ($r=0.358^*$), productive tillers per plant ($r=0.490^*$), spikelets per panicle ($r=0.422^*$), spikelet fertility ($r=0.456^*$), harvest index ($r=0.369^*$) and 1000 grain weight ($r=0.553^{**}$) indicated that while practicing selection these traits has to be give prime importance in early generations (Table 4). Similar results reported by Hari Rama Krishnan *et al.*, 2006 and Lokhendra singh *et al.*, 2008)

The path analysis revealed that the highest direct effect on grain yield per plant is exhibited by biomass (2.7658) followed by panicle weight (2.2292) and days to maturity (0.1761). The traits with significant and positive correlation can be used for selection to improve the grain yield and the traits with highest direct effects also helpful in improving grain yield through the selection for those traits (Karim *et al.*, 2007; Debnath *et al.*, 2015).

Validation of trait specific markers

Out of 500 markers, 78 were polymorphic on 3.5% percent agarose gel. On an average, 14.18 per cent polymorphism was obtained. Similar type of results were obtained by Shishido *et al.*, 2006 indicating use of these polymorphic markers for identification and selection of good progeny plant types for associated traits and also for fingerprinting of parents. Among the polymorphic markers twenty trait specific SSR markers were used for validation, among the twenty marker chosen for the study, only six were found to be significantly associated with yield traits. Details of SSR markers association found for different characters are furnished in table.5 . It has been found that RM495 is associated with yield contributing traits spikelet fertility (%) and plant height at maturity (cm) scores with phenotypic data under field conditions showed significance at 5% level. RM 242 was found to be significantly associated with grain yield at 5% level of significance. RM 152 was found to be significantly associated with spikelet fertility (%) at 5% level of significance. RM 263 found to be associated with grain yield at 5% level of significance. Only one marker RM 248 showed significant association with days to fifty percent flowering at 5% level of significance. RM 596 was found to be associated with grain yield at 5 % level of significance similar results reported by Kanagaraj *et al.* (2010).

The percentage contribution of the significantly associated markers to total phenotypic variance (R^2) varied from 4.45 in case of RM 596 for grain yield at harvest to 19.73 in case of RM 495 spikelet fertility. In case of RM 242 phenotypic variance

explained is (R^2) is 15.31 for grain yield, RM 152 explained phenotypic variance of 12.06 (R^2) for the trait spikelet fertility, RM 248 explained days to 50% flowering phenotypic variance (R^2) is 10.23. Low R^2 values indicates large environmental component of variation. This low R^2 is due to large QTLs are controlling the traits. The identified polymorphic markers will be further utilized for development of linkage map of cross IM90 X IR 64. The genotypes carrying number of positive alleles were given in table 6. These genotypes could be selected and advanced to next generation as they were carrying alleles similar to parent IM90 which is high yielding genotype.

REFERENCES

- Bidhan, R., Hossain M., Hossain F. and Roy A. 2011.** Genetic variability in yield components of rice (*Oryza sativa* L.), *Envt. and Ecol.* **19**: 186-189.
- Collard, B. C. Y., Jahufer, M. Z. Z., Brouwer, J. B. and Pang, E. C. K. 2005.** An introduction to markers, quantitative trait loci (QTL) mapping and marker-assisted selection for crop improvement: The basic concepts. *Euphytica.* **142**: 169-196.
- Debnath, K., Das, B., Sikder, S. and Sarkar K. K. 2015.** Assessment of genetic variability character association and path coefficient for yield and its component characters in rice. *The Ecoscan.* **9(1&2)**: 455-459.
- Dellaporta, S. L., Wood, J. and Hicks, J. B. 1983.** A plant DNA-mini preparation version II, *Plant Molecular Biology Reporter.* **1**:19-21.
- Edwards, M. D., Stubber, C. W. and Wendel, J. F. 1987.** Molecular marker facilitated investigations of quantitative trait loci in Maize. Number, genomic distributions and type of gene action. *Genetics.* **116**: 113-125.
- FAOSTAT. 2013.** Statistical data base, Food and Agricultural Organization (FAO) of the United Nations, Rome, <http://www.fao.org>.
- FarhadKahani and ShailajaHittalmani. 2015.** Genetic Analysis and Traits Association in F_2 Intervarietal Populations in Rice under Aerobic Condition, *J. Rice Res.* **3(152)**: 125-132.
- Federer, W. T. 1956.** Augmented (or hoonuiaku) designs. *Hawaii. Plant Research.* **2**: 191-208.
- Gandhi, V. R., Hittamani, S. and Shivamurthy, M. 2011b.** Performance and adaption of Direct seeded aerobic rice variety MAS26 (Onasiri) in Southern Karnataka. *J. EXT. Edu.* **23(3)**: 4684-4690.
- Gandhi, V. R., Shivamurthy, M., Rudresh, N.S. and Hittamani, S. 2011a.** Performance and adaption of new aerobic rice variety MAS946-1 (Sharada) in southern Karnataka. *Karnataka J Agric. Sci.* **25 (1)**:5-8.
- Gangashetty, P., Salimath, P. and Hanamaratti, N. 2013.** Genetic variability studies in genetically diverse non-basmati local aromatic genotypes of rice (*Oryza sativa* L.), *Rice Genomics and Genetics.* **4(2)**:8-12
- Ghosh, S. C. and Sharma, D. 2012.** Genetic parameters of agro morpho- physiological traits in rice (*Oryzasativa* L.), *Electronic J. of Pl. Breed.* **3(1)**: 711-714.
- Hariramkrishnan, S., Anandakumar, C. R., Saravanan, S., and Malini, N. 2006.** Association Analysis of Some Yield Traits in Rice (*Oryza sativa* L.), *J. Applied Sci. Res.* **2**: 402-404.
- Idris, A. E., Justin, F. J., Dagash, Y. M. I., and Abuali, A. I. 2012.** Genetic variability and inter relationship between yield and yield components in some rice genotypes. *American. J. Experi Agric.* **2(2)**: 233-239.
- Kanagaraj P., Prince, K. S. J., Sheeba, J. A., Biji, K. R., Paul, S. B., Senthil, A. and Babu, R. C., 2010.** Microsatellite markers linked to drought resistance in rice (*Oryza sativa* L.). *Curr. Sci.* **98(6)**: 836-839.

- Karim, D., Sarkar, U., Siddique, M. N. A., KhalequeMiah, M. A. and Hasnat, M. Z. 2007. Variability and Genetic Parameter Analysis in Aromatic Rice. *Int. J. Sustain. Crop Prod.* **2(5)**:15-18.
- Khalid, A. O., Ahmed, M. M., Farhan, A., Zheng, Y. and Qiu, F. 2012. Genetic variability for yield and related attributes of upland rice genotypes in semiarid zone (Sudan). *African J. Agric Res.* **7(33)**: 4613-4619.
- Kumar, A., Verulkar, S., Dixit, S., Chauhan, B., Bernier, J., Venuprasad, R., Zhao, D. and Shrivastava, M. N. 2009. Yield and yield-attributing traits of rice (*Oryza sativa* L.) under lowland drought and suitability of early vigor as a selection criterion. *Field Crops Res.* **114(1)**: 99-107.
- Kumar, R. G., Suresh Babu, Satish Kumar R. and Sandhya. 2015. Genetic variability and divergence studies in elite rice genotypes for biometrical traits. *The Eoscan.* **9(1&2)**: 209-212.
- Lafitte, H., Z. Li., Vijayakumar, C., Gao, Y., Shi, Y., Xu, J., B. Fu, S. Yu, Ali, A. and Domingo, J. 2006. Improvement of rice drought tolerance through backcross breeding: evaluation of donors and selection in drought nurseries. *Field Crop Res.* **97**: 77-86.
- Laxuman. 2010. Molecular mapping and tagging of quantitative Trait LOCI (QTLs) for Morphological physiological and yield traits in rice (*Oryza sativa* L.). *Ph.D. Thesis*, University of Agricultural Sciences, Dharwad.
- Lokhendrasingh, Daya Ram and BrahamSingh. 2008. Genetic variability and character association in paddy. *Society of Sci. Dev. In Agric. and Tech.* **3(2)**: 209-210.
- Manjappa, Uday, G. And Hittalmani, S. 2014. Association analysis of drought and yield related traits in F₂ population of moroberekan/ir64 rice cross under aerobic condition. *TJPRC.* **4(2)**: 79-88.
- Marri, P. R., Sarla, N., Reddy, V. L. N. and Siddique, E. A. 2005. Identification and mapping of yield and yield related QTLs from an Indian accession of *Oryzaru fipogon*. *BMC Genet.* UK. **6**: 33-40.
- NareshBabu, N., Shivakumar, N and ShailajaHittalmani. 2010. Pollen fertility Vs Spikelet fertility in F₂ of a CMS based hybrids in rice (*Oryza sativa* L.) under aerobic condition. *Electronic J Pl Breed.* **1(4)**: 789-793.
- Ogunbayo, S. A., Sie, M., Ojo, D. K., Akinwale, M. G., Toulou, B., Shittu, A., Idehen, E. O., Popoola, A. R. and Gregorio, G. B. 2014. Genetic variation and heritability of yield and yield related traits in promising rice genotypes (*Oryzasativa* L.), *J. Plant breeding and crop science.* **6(11)**: 153-159.
- Paikhomba, N., Kumar, A., Chaurasia, A. K. and Rai, P. K. 2014. Assessment of Genetic Parameters for Yield and Yield Components in Hybrid Rice and Parents. *J. Rice Res.* **2**: 117.
- Panse, V. G. and Sukhatme, P. V. 1967. Statistical methods for agricultural workers, ICAR, New Delhi, P.360.
- Powell, W., Machray, G. C. and Provan, J. 1996. Polymorphism as revealed by simple sequence repeats. *TIPS.* **1**: 215-222.
- Prajapati, M. K., Singh, C. M. Babu, S., Lavanya, P. C. and Jadav, P. 2011. Genetic parameters for grain yield and its component characters in rice. *Electronic J of Pl. Breed.* **2(2)**: 235-238.
- Price, A. H., Cairns, J. E., Horton, P., Jones, H. G. and Griffiths, H. 2002. Linking drought resistance mechanisms to drought avoidance in upland rice using a QTL approach: progress and new opportunities to integrate stomatal and mesophyll responses. *J. of Exp. Bot.* **53**: 989-1004.
- Raghavendra, P and Hittalmani, S. 2015. Genetic parameters of two BC₂F₁ populations for development of superior male sterile lines pertaining to morpho-floral traits for aerobic rice (*Oryzasativa* L.), *SAARC J. Agri.* **13(2)**: 198-213.
- Robinson, H. F., Comstock, R. E. and Harney, P. H. 1949. Estimates of heritability and degree in corn. *Agron. J.* **41**: 353-359.
- Shantavva, G., Mahadevu, P., VeeraghantiSmita, S., Suresh, K. and Hittalmani, S. 2014. Variability Studies for Yield and its Components in Rice (*Oryzasativa* L.), *Trends in Biosciences.* **7(10)**: 961- 963.
- Shailaja Hittalmani and Shivashankar, G. 1987. Out crossing on male sterile lines of rice. *Mysore J. Agricultural Sciences.* **21**: 158-160.
- Shishido, R., Kikuchi, M., Nomura, K. and Ikehashi, H. 2006. Evaluation of genetic diversity of wild rice (*Oryzarufipogon*Griff.) in Myanmar using simple sequence repeats (SSRs). *Genet. Resources and Crop Evol.* **53**: 179-186.
- Standard Evaluation System (SES) for Rice. 1996. Prepared by INGER – Genetic Resources Center, International Rice Research Institute, Manila, Philippines. PP.1-39.
- Venuprasad, R., Impa, S., Bernier, J., Serraj, R., Kumar, A., and Gowda, R. P. V. 2009. The large-effect drought-resistance QTL *qtl12.1* increases water uptake in upland rice. *Field Crops Res.* **110**: 39-46.
- Xing, Y. Z., Tan, Y. F., Hua, J. P., Sun, X. I., Xu, C. G. and Zhang, Q. 2002. Characterization of the main effects, epistatic effects and their environmental interactions of QTLs on the genetic basis of field traits in rice. *Theor. Appl. Genet.* **105**: 248-257.
- Yue, B., Xiong, L., Xue, W., Xing, Y., Luo, L. and Xu, C. 2005. Genetic analysis for drought resistance of rice at reproductive stage in field with different types of soil. *Theor. Appl. Genet.* **111**: 1127-1136.
- Yue, B., Xue, W., Xiong, L., Yu, X., Luo, L., Cui, K., Jin, D., Xing, Y. And Zhang, Q. 2006. Genetic Basis of Drought Resistance at Reproductive Stage in Rice: Separation of Drought Tolerance From Drought Avoidance. *Genetics.* **172**: 1213-1228.
- Zhang, J., Zheng, H. G., Aarti, A., Pantuwan, G., Nguyen, T. T., Tripathy, J. N., Sarial, A. K., Robin, S., Babu, R. C., Nguyen, B. D., Sarkarung, S., Blum, A. and Nguyen, H. T. 2001. Locating genomic regions associated with components of drought resistance in rice: comparative mapping within and across species. *Theor. Appl. Genet.* **103**: 19-29.