



ISSN: 0974 - 0376

*The Ecoscan* : Special issue, Vol. VII: 07-12: 2015  
AN INTERNATIONAL QUARTERLY JOURNAL OF ENVIRONMENTAL SCIENCES  
[www.theecoscan.in](http://www.theecoscan.in)

## IMPACT OF WEATHER FACTORS ON THE INCIDENCE OF MAJOR INSECT PESTS OF TOMATO (*SOLANUM LYCOPERSICON* L.) CV. H-86 (KASHI VISHESH)

Rishikesh Mandloi *et al.*,

### KEYWORDS

Tomato  
Seasonal incidence  
Jassid  
Aphid  
White fly  
Thrips  
Fruit borer

Proceedings of National Conference on  
Harmony with Nature in Context of  
Bioresources and Environmental Health  
(HARMONY - 2015)  
November 23 - 25, 2015, Aurangabad,  
organized by  
Department of Zoology,  
Dr. Babasaheb Ambedkar Marathwada University  
Aurangabad (Maharashtra) 431 004  
in association with  
NATIONAL ENVIRONMENTALISTS ASSOCIATION, INDIA  
[www.neaindia.org](http://www.neaindia.org)



RISHIKESHMANDLOI\* RAJESH PACHORI, AMIT KUMAR SHARMA, MONI THOMAS AND A. S. THAKUR

Department of Entomology,

College of Agriculture Jabalpur Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur - 482 004 (Madhya Pradesh) INDIA

e-mail: rishikeshmandloi1@gmail.com

## ABSTRACT

Major activity period of *Aphis gossypii* Glover was observed from October 2012 to March 2013 with two distinct peaks during 7<sup>th</sup> and 11<sup>th</sup> SW (standard week) (11.22 and 11.66 aphid/ 6 leaves) and *Liriomyza trifolii* Burgess was observed from October 2012 to March 2013 with three distinct peaks during 10<sup>th</sup>, 11<sup>th</sup> and 12<sup>th</sup> SW (44.56%, 45.95% and 44.02%) respectively. While, *Bemisia tabaci* Genn was appeared November 2012 to March 2013 with two distinct peaks during 7<sup>th</sup> and 9<sup>th</sup> SW (9.84 and 11.85 flies/10cm twig). *Amrasca devastans* Ishida, and *Scirtothrips dorsalis* Hood were observed during November 2012 to March 2013 with two distinct peaks 9<sup>th</sup> and 11<sup>th</sup> SW (9.26 and 9.15 jassid/6 leaves) and 7<sup>th</sup> and 9<sup>th</sup> SW (2.08 and 1.85 thrips/ 6 leaves) respectively. *Helicoverpa armigera* Hub. of observed from November 2012 to March 2013 with two distinct peaks 11<sup>th</sup> and 12<sup>th</sup> SW (6.02 and 6.11 larvae/plant). Analysis of correlation coefficient between abiotic factors (weather parameters) and the major insect pests of tomato showed that population of thrips had a significant positive correlation with evening relative humidity (R.H.) (0.456\*), while fruit borer had a significant positive correlation with rainy days (0.428\*).

## INTRODUCTION

Tomato is the most important vegetable grown widely both for fresh marketing and processing. India is next only to the China in area and production of vegetables. In Madhya Pradesh, it is grown in 60.84 thousand hectare with the annual production of 1484.5 thousand metric tons and productivity of 24.40 metric tons hec<sup>-1</sup> (Anonymous). The production and quality of tomato fruits are considerably affected by array of insect pests infesting at different stages of crop growth. The key insect-pests of tomato includes Aphid (*Aphis gossypii* Glover), Jassid (*Amrasca devastans* Ishida), White fly (*Bemisia tabaci* Genn.), Leaf miner (*Liriomyza trifolii* Burgess), Thrips (*Scirtothrips dorsalis* Hood) and Fruit borer (*Helicoverpa armigera* Hub.) Choudhri *et al.* (2001), Reddy and Kumar (2004). Kharpuse *et al.* (2005) revealed that the fruit borer population peaked (26 larvae/10 plants) uniformly distributed. According to Mathur *et al.* (2012) Pest abundance and distribution changes with abiotic factors and therefore meteorological parameters play a pivotal role in the biology of any pest. Subba *et al.* (2014) revealed that the lower population level was recorded during 38<sup>th</sup> standard week to 9<sup>th</sup> SW and 19<sup>th</sup> SW to 20<sup>th</sup> SW. Higher population level was maintained during 10<sup>th</sup> SW to 20<sup>th</sup> SW and peak population (1.11/plant) was recorded on 23<sup>rd</sup> SW. Abiotic conditions such as relative humidity (maximum, minimum and average) had significant negative influence on *Phyllotreta spp.* while non-significant negative correlation with weekly total rainfall. In case of temperature (maximum, minimum, and average), non-significant positive correlation was observed. Temperature influencing the rate of growth and development of insect. However, relative humidity, rainfall, wind speed and temperature are the chief weather parameters that largely direct the activity of a given species of insect. The interaction between pest activity and abiotic factors helps in deriving at predictive models that aids in forecast of pest incidence. The present study was aimed to know the seasonal occurrence of different insect pest and the effect of abiotic factors on their population in Jabalpur region of M.P.

## MATERIALS AND METHODS

Present experiment was conducted at Vegetable Research Farm, Department of Horticulture, College of Agriculture, JNKVV, Jabalpur (M.P.) during Rabi 2012-13. Seasonal incidence of major insect pests *viz.*, Aphid (*Aphis gossypii* Glover), Jassid (*Amrasca devastans* Ishida), White fly (*Bemisia tabaci* Genn.), Leaf miner (*Liriomyza trifolii* Burgess), Thrips (*Scirtothrips dorsalis* Hood) and Fruit borer (*Helicoverpa armigera* Hub.) were studied during the season. The Observations on major insect pests were recorded on 25 randomly selected plants of the crop in a standard week (7 day interval) from transplanting to till the availability of insects or maturity of the crop. Kumar (2008) as per the method proposed by. Observations on sucking and miner insect pests population (nymphs and adults) of Aphid, Jassid, Thrips, Leaf miner and whitefly were taken on 6 leaves plant<sup>-1</sup> *viz.* (two upper, middle and lower plant canopies) and in case of fruit borer larval population was recorded on plant<sup>-1</sup> basis. Observations were statistically analysed in correlation analysis described by Panse and Sukhatme (1963). The impact of abiotic factors on population

\*Corresponding author

dynamics of major insect pest species of tomato Infested fruit was recorded and percentage fruit damaged were worked out.

## RESULTS AND DISCUSSION

Incidence of major insect pests of tomato during *Rabi* 2012-13 and Correlation coefficient of insect population with abiotic factors (weather parameters) as follows.

### Aphid (*A. gossypii* Glover)

Aphid was first recorded during 43<sup>th</sup>SW on tomato. The activity period of aphid was observed from 43<sup>th</sup> to 12<sup>th</sup> SW (22 Oct

2012 to 25 March 2013). With two distinct peaks during 7<sup>th</sup> and 11<sup>th</sup> SW (11.22 and 11.66 aphid/6 leaves). The highest peak was observed in 11<sup>th</sup> SW (11.66 aphid/6 leaves) during this period was maximum temperature (31.6°C) and minimum/temperature (14.7°C), whereas morning and evening relative humidity and rainy days (no.) were 84% and 39% and 02 days respectively. Thus, the pest was present throughout the growing stage of the crop. It caused leaf curl, defoliation, stunted growth of plants and development of sooty mould on the leaves. Present findings are in accordance with those of Chaudhuri *et al.* (2001), Reddy and Kumar (2004), Umesh and Onukwu (2005) and Mandal (2012). These authors also


**Table 1: Weekly meteorological data and incidence of major insect pests on tomato during *Rabi* 2012-13**

Standard Week nos.	Temperature (°C)		Relative humidity (%)		Rainy days (no.)
	Max	Min	Morning	Evening	
43	21.6	15.2	82	35	0
44	27.9	15.6	91	51	0
45	28.2	14.6	88	30	0
46	28.6	10.0	85	28	0
47	27.9	10.6	88	29	0
48	28.4	11.5	83	33	0
49	28.7	10.6	85	26	0
50	29.0	14.0	92	41	0
51	25.3	7.1	88	29	0
52	23.8	5.0	90	30	0
1	23.3	7.2	87	42	0
2	23.0	5.2	87	32	0
3	26.7	10.1	84	36	0
4	21.4	5.0	86	36	0
5	24.6	7.4	91	36	0
6	25.9	11.3	88	49	0
7	25.2	13.0	91	60	0
8	25.0	11.0	93	49	3
9	28.0	9.2	87	37	1
10	30.7	9.9	83	28	0
11	31.6	14.7	84	39	2
12	33.4	16.2	77	28	0

Aphid Per six leaves	Jassid	Thrips	Whitefly population /10 cm twig	Leaf miner infestation (%)	Fruit borer larval population/plant
0.07	0.00	0.00	0.00	0.14	0.00
0.06	0.00	0.00	0.00	1.22	0.00
0.07	0.20	0.00	0.76	4.70	0.00
0.55	0.56	0.05	0.65	8.33	0.00
1.58	1.77	0.30	0.54	12.59	0.00
1.61	2.73	0.51	1.32	13.55	0.51
2.42	2.42	0.59	1.76	13.46	0.70
1.96	1.11	0.68	0.91	14.70	0.89
3.35	3.39	1.01	3.66	16.29	1.04
3.37	4.01	1.12	4.00	18.62	1.54
4.08	3.87	1.32	4.44	21.82	1.77
5.09	4.74	1.15	5.11	26.28	1.91
6.12	5.24	1.31	5.85	28.40	2.09
7.30	6.21	1.15	4.66	32.43	2.67
8.62	6.54	1.46	6.33	34.29	3.12
10.30	8.00	1.83	7.41	33.18	3.84
11.22	7.99	2.08	9.84	32.74	4.66
4.69	8.00	1.42	4.51	39.04	4.90
10.65	9.26	1.85	11.85	41.80	5.29
8.42	8.59	1.50	9.10	44.56	5.50
11.66	9.15	1.58	8.44	45.95	6.02
7.66	7.00	0.00	6.09	44.02	6.11

**Table 2: Correlation coefficient of insect population with Meteorological parameter**

Weather Parameter	Aphid r	Jassid r	Whitefly r	Leaf miner r	Thrips r	Fruit borer r
Maximum Temp. (°C)	0.116	0.115	0.106	0.231	-0.226	0.224
Minimum Temp.(°C)	0.121	0.189	-0.193	-0.155	-0.421	0.289
Morning RH (%)	0.066	0.060	-0.060	-0.148	0.317	-0.086
Evening RH (%)	0.336	0.280	0.275	0.161	0.456*	0.414
Rainy days	0.379	0.389	0.416	0.386	0.333	0.428*

**Figure 1: Incidence of major insect pests Aphid, Jassid and Thrips on tomato during Rabi 2012-13**

reported that the, *A. gossypii* to be an important sucking pest of tomato and was present throughout the growing period of the crop. Hath and Das (2004) recorded maximum *A. gossypii* population on tomato during the first week of March.

Correlation coefficient between various weather parameters and *A. gossypii* population were found no significant.


#### Jassid (*A. devastans* Ishida)

*A. devastans* population was first recorded during 45<sup>th</sup> SW on tomato. The activity period of *A. devastans* was observed from 45<sup>th</sup> to 12<sup>th</sup> SW (05 November 2012 to 25 March 2013). With two distinct peaks during 9<sup>th</sup> and 11<sup>th</sup> SW (9.26 and 9.15 jassid/ 6 leaves). The highest peak was observed 9<sup>th</sup> SW (9.26 jassid/ 6 leaves). During this period maximum and minimum temperature were 28.0°C and 9.2°C respectively, whereas morning and evening relative humidity and rainy days (no.) were 87% and 37% and 01 days respectively. The pest was present throughout the growing stage of the crop, and caused leaf curl leading to stunted growth of plants. It was one of the major sucking pests of tomato. Present observations were more or less similar with the results of earlier workers Kumar (2008) reported that the *A. devastans* appeared in the 1<sup>st</sup> week of January.

Correlation coefficient of *A. devastans* population exhibited non significant relationship with all the weather parameter.

#### White fly (*B. tabaci* Genn.)

*B. tabaci* was first appeared during 45<sup>th</sup> SW on tomato plants foliage canopy. The activity period of *B. tabaci* was observed from 45<sup>th</sup> to 12<sup>th</sup> SW (05 November 2012 to 25 March 2013) with two distinct peak during 7<sup>th</sup> and 9<sup>th</sup> SW (9.84 and 11.85 flies/ 10cm twig). The highest peak was observed 9<sup>th</sup> SW (11.85

**Figure 2: Incidence of major insect pest White Fly, Leaf miner and Fruit borer on Tomato during Rabi 2012 -13**

flies/10 cm twig). During this period maximum and minimum temperatures were 28.0°C and 9.2°C respectively, whereas morning and evening relative humidity and rainy days (no.) were 87% and 37% and 01 days respectively. *B. tabaci* population ranged from (0.0 to 11.85 flies/10 cm twig) with an overall mean performance of (4.42 flies/10 cm twig). Similar findings have been reported by Chaudhuri *et al.* (2001), Reddy and Kumar (2004), and Mandal (2012). They also reported, *B. tabaci* to be an important sucking pest of tomato and were present throughout the growing period of the crop. Kharpuse (2005) and Chaudhuri *et al.* (2001) found that the *B. tabaci* population peaked (13 files/10 cm twig) during first week of March.

Correlation coefficient between various weather parameters and *B. tabaci* population expressed no significant relationship.

#### Leaf miner (*L. trifolii* Burgess)

*L. trifolii* was first appeared 43<sup>th</sup> SW on tomato plants and continued till the harvesting. Increasing trend was observed in this case with crop growth stages. The activity was observed 43<sup>rd</sup> to 12<sup>th</sup> SW (22 Oct 2012 to 25 March 2013) with three distinct peaks during 10<sup>th</sup>, 11<sup>th</sup> and 12<sup>th</sup> SW (44.56%, 45.95% and 44.02%) and maximum leaf infestation was recorded on 11<sup>th</sup> SW (45.95%). During this period was maximum temperature (31.6°C) and minimum temperature (14.7°C), whereas morning and evening relative humidity and rainy days (no.) were 84% and 39% and 02 days respectively. The pest was present throughout the growing stage of the crop and mined the leaves. Present findings are in accordance with those of Marcano and Issa (2000), Chaudhuri *et al.* (2001), Asalatha (2002) and Reddy and Kumar (2004). They recorded



Plate 1: Visual symptoms of incidence of major insects on tomato at Jabalpur. (a) *H. armigera* on tomato fruit, (b) *A. devastans* on tomato leaf, (c) *B. tabaci* on tomato leaf, (d) *A. gossypii* on tomato leaves and (e) *L. trifolii* on tomato leaves

*Liriomyza* spp. as major pest in tomato. Kharpuse (2005) revealed that the maximum (76.67%) leaf infestation by *L. trifolii* was recorded at middle of the March.

Correlation coefficient between various weather parameters and *L. trifolii* tunnel was found no significant relationship.

#### Thrips (*Scirtothrips dorsalis* Hood)

*S. dorsalis* appeared on 46<sup>th</sup> SW on tomato plants. The activity period of *S. dorsalis* was observed from 46<sup>th</sup> to 11<sup>th</sup> SW (12 November 2012 to 18 November 2013) and maximum peak population was appeared in 7<sup>th</sup> SW (2.08 %). During this period was maximum temperature (25.2°C) and minimum temperature (13.2°C), whereas morning and evening relative humidity and rainy days (no.) were 91% and 60% and 00 days respectively. Present findings are in accordance with those of Rajkumar *et al.* (2003) and Goncalves *et al.* (2004).

Correlation coefficient various weather parameters and *S. dorsalis* population were found non significant, except evening relative humidity which exhibited a significant and positive

correlation (0.456\*).

#### Fruit borer (*Helicoverpa armigera* Hub.)

The larvae of *H. armigera* were first observed on tomato crop during 48<sup>th</sup> SW. The activity period of *H. armigera* was observed from 48<sup>th</sup> to 12<sup>th</sup> SW (26 November 2012 to 25 March 2013) and gradually increased till the crop was harvested. Maximum peak of larval fruit infestation was appeared in 12<sup>th</sup> SW (6.11 larvae/plant<sup>-1</sup>), during this period was maximum temperature (33.4°C) and minimum temperature (16.2°C), whereas morning and evening relative humidity and rainy days (no.) were 77% and 28% and 00 days respectively. The pest was present during the entire reproductive stage of the crop and caused circular or irregular holes on the surface of the fruit and bore inside it. Present findings are in accordance with those of Rudenko *et al.* (2001), Chaudhuri *et al.* (2001), Reddy and Kumar (2004) and Mandal (2012). They all reported that *H. armigera* had been major insect pests of tomato. Meena L.K. *et al.* (2014) revealed that the incidence of *H. armigera* on tomato was recorded at regular weekly interval and correlated

with the corresponding meteorological data the incidence of *H. armigera* was also observed throughout the cropping season from 18.10.2011 to 25.02.2012. Kharpuse (2005) revealed that the fruit borer population peaked (26 larvae/10 plant) in the third week of March.

Correlation coefficient various weather parameters and *S. dorsalis* population were found non significant, except rainy day which exhibited a significant and positive correlation (0.428\*).

## ACKNOWLEDGMENT

Thanks to most esteemed guide and my advisory Dr. Rajesh Pachori (Professor), Dr. A.S. Thakur (Scientist), Dr. Moni Thomas (Senior Scientist), and especially thanks to Dr. A.K. Sharma (Assistant Professor) for his cheerfully advice during thesis period, Department of Entomology, College of Agriculture, JNKVV, Jabalpur for his most valuable and inspiring guidance, close supervision, constant encouragement and constructive criticism coupled with valuable suggestions during the course of this investigation and in preparation of this manuscript.

## REFERENCES

- Asalatha, R. 2002.** Seasonal activity and bio-efficacy of some eco-friendly insecticides against the serpentine leaf miner *Liriomyzatrifolii*. M.Sc. (Ag.) Thesis, JNKVV, Jabalpur.
- Chaudhuri, N. and Senapati, S. K. 2001.** Evaluation of pesticides from different origin-synthetic and biological, against pest complex of tomato under terai region of West Bengal. *Haryana J. Horti. Sci.* **30(3/4)**: 274-277.
- Goncalves, P. A. S., Werner, H. and Debarba, J. F. 2004.** Evaluation of biofertilizers, plant extracts, and some alternative substances to manage onion thrips in organic agriculture system. [Portuguese]. *Horti. Brasileira.* **22(3)**: 659-662.
- Hath, T. K. and Das, B. R. 2004.** Incidence of insect pests in late planted tomato under terai agroecology of West Bengal. *Environ. Ecol.* **22(1)**: 136-140.
- Kharpuse, Y. K. 2005.** Studies on seasonal incidence and role of botanical against major insect pests of tomato (*L. esculentum* M.). M.Sc. (Ag.) (Ent.) Thesis submitted to J.N.K.V.V., Jabalpur (M.P.). pp. 1-53.
- Kumar, K. Indra 2008.** Studies on insect pest complex of tomato (*Lycopersicon esculentum* Mill.) and management of fruit borer, *Helicoverpa armigera* Hub. with chemicals. M.Sc. (Ag.) (Ent) Thesis submitted to J.N.K.V.V., Jabalpur (M.P.). pp. 1-91.
- Mandal, S. K. 2012.** Bio-efficacy of cyazypyr 10% OD, a new anthranilic diamide insecticide, against the insect pests of tomato and its impact on natural enemies and crop health. *Acta Phytopathologica et Entomologica Hungarica.* **47(2)**: 233-249.
- Mathur Anjali\*, Singh N. P., Meena Mahesh<sup>1</sup> and Singh Swaroop 2012.** Seasonal incidence and effect of abiotic factors on population dynamics of major insect pests on brinjal crop. *J. Environ. Res. Develop* vol. 7 no. 1a,
- Meena, L. K. and Bairwa, B. 2014.** Influence of abiotic and biotic factors on the incidence of major insect pest of tomato. *The Ecoscan* pp. **(3&4)**: 309-313
- Marcano, R. and Issa, S. 2000.** Spatial and vertical distribution of *Liriomyzatrifolii* on tomato. *Boletin de Entomologica venezolana* **8(1)**: 115-122.
- Panse, V. C. and Sukhatme, P. V. 1963.** Statistical methods for agricultural workers. *ICAR Publications*, New Delhi.
- Rudenko, N. E., Zubanov, A. P., Sherbinin, B. H. and Chalenko, V. V. 2001.** Pest control by the pneumatic method. *Zeschita Rastenii.* pp. 10-16.
- Rajkumar, A. J. Kurian, P. S. Backiyarani, S. and Murugan, M. 2003.** Evaluation of biorationals against thrips (*Sciothrips cardamomi* Ramk.) and shoot and capsule borer (*Conogethes punctiferalis* Guen.) in cardamom. *J. Spices and Aromatic Crops.* **11(2)**: 132-134.
- Reddy, N. A. and Kumar, C. T. A. 2004.** Insect pests of tomato, *Lycopersicon esculentum* Mill. in eastern dry zone of Karnataka. *Insect Environ.* **10(1)**: 40-42.
- Subba, B., Ghosh, S. K., Ravikumar, K. and Chhetri, B. 2014.** Seasonal incidence of flea beetle (*Phyllotreta* spp.) infesting tomato (*Lycopersicon esculentum* L.) and their sustainable management. *The Ecoscan, Ppvol.* **6**: 175-180
- Umesh, V. C. and Onukwu, D. 2005.** Development of environmentally-friendly tomato insect pest control options under tropical conditions. *J. Vege. Sci.* **11(3)**: 73-84.