

DYNAMICS OF SPECIES COMPOSITION OF STEM BORERS IN RICE CROP

AMIT KUMAR*, A. K. MISRA, PAIDI SATYANARAYANA AND SHANTA KUMAR CHOUDHARY¹

Department of Entomology,
Rajendra Agricultural University, Pusa, Samastipur - 848 125, INDIA
¹Research Associate, GKMS, Agro Meteorology,
Rajendra Agricultural University, Pusa, Samastipur - 848 125, INDIA
e-mail: amitkumarpusa1@gmail.com

INTRODUCTION

Rice (*Oryza sativa*) belonging to the family Graminae, is one of the most important staple food crops not only in India but the world too. Rice is grown in all the continents except Antarctica, occupying 158 million hectare in 111 countries in the world (Agricultural statistics at a glance, 2011), out of which Asia accounts for 90 per cent and America, Australia, Africa and Europe cover the rest 10 per cent (Ampong-Nayarko and De Datta, 1991, Roy *et al.*, 2013 and Saxena and Murty 2014) with a production of 685 million tonnes and productivity of 4328kg/ha (Agricultural statistics at a glance, 2011).

Rice is grown in both *kharif* and *rabi* season under diverse ecological and climatic conditions apart from socio-economic diversities of the state. The unholy triple alliance of pests (insects, diseases and weeds) act as a great impediment in achieving desired level of rice production (Gupta *et al.*, 2002, Balakrishna and Satyanarayana 2013 and Manikandan *et al.*, 2014). Intensive use high yielding varieties, sequential cropping and indiscriminate use of insecticides have resulted in various insect pest problems in rice crop. Among various depressing factors, biotic stress as insect pest infestation is the most crucial factor due to which rice production is unpredictable. In India, approximately 100 insect pests have been reported as pests of rice and 20 of these are considered to be major pests causing 30 per cent yield loss from seedling to maturity (Cramer, 1967; Pathak and Dhaliwal, 1981 and Atwal and Dhaliwal, 2005). All together 21 species of lepidopteran stem borers have been recorded as rice pests throughout the world. Of these, 8 species are known to occur in India (Rao, 1965 and Pathak, 1975). Of various lepidopteran insect pests attacking on rice, yellow stem borer, *Scirpophaga incertulas* Walker, white stem borer, *Scirpophaga innotata* (Walker), dark headed borer, *Chilo polychrysus* (Meyrick) and pink stem borer, *Sesamia inferens* (Walker) are economically important. They damage most of the crop and harm the productivity of Rice. The newly emerged larva enters into the stem for feeding on inner tissues at vegetative and reproductive stage of the crop. As a result of their feeding inside the stem around the nodes, central leaf whorl remains unfold, turn brownish, dry up and easily be pulled out, while lower leaves remain green and healthy. This condition is known as deadheart (DH). The affected tillers do not produce panicles. If infestation continues to the ripening stage of the crop then plants bear panicles without grains (chaffy ears). This condition is known as white earhead (WEH). Keeping the harmful nature of these stem borer in view, the present investigation is undertaken in agro-climatic zone-I of North Bihar.

MATERIALS AND METHODS

In order to study the dynamics of pest species composition of stem borers of rice viz. yellow stem borer, *Scirpophaga incertulas* (Walker), pink stem borer, *Sesamia*

ABSTRACT

As the stem borers are harmful insect for rice crop it is very important to know the dominant species population to management of these insect. In order to determine the population dynamics of harmful stem borer to develop ecological and economic viable strategies, a field experiment was conducted at the Research Farm, R.A.U., Pusa and in the Laboratory, Department of Entomology, R.A.U., Pusa. Results pertaining to the dynamics of species composition of stem borers of rice reveal that all the four species of stem borer of rice viz. yellow stem borer, *Scirpophaga incertulas* (Walker), pink stem borer, *Sesamia inferens* (Walker), white stem borer, *Scirpophaga innotata* (Walker) and dark headed striped borer, *Chilo polychrysus* (Meyrick) were prevalent during the crop season. The population of yellow stem borer was found to be dominant over other species of stem borer and recorded 87.00% to 93.00% at 30 DAT during two *kharif* seasons respectively. The population was 82.50% to 89.00% at 50 DAT and 85.00% to 91.50% at 90 DAT.

KEY WORDS

Chilo polychrysus (Meyrick)
Rice
Scirpophaga incertulas (Walker)
Scirpophaga innotata (Walker)

Received : 11.09.2015
Revised : 20.03.2016
Accepted : 07.05.2016

*Corresponding author

inferens (Walker), white stem borer, *Scirpophaga innotata* (Walker) and dark headed striped borer, *Chilo polychrysus* (Meyrick) in North Bihar condition, a field trial was conducted at research farm, R.A.U., Pusa, Samastipur, Bihar during *Kharif*, 2010 and *Kharif*, 2011. For this purpose, field preparation was done by ploughing once by tractor drawn disc plough and second ploughing was done by mould board plough. A well established susceptible variety Rajendra Kasturi was transplanted with 15cm (plant to plant) x 20cm (row to row) spacing in 100 sq.m plot area. The entire plot was quarterised into equal sized blocks and demarcated with bunds (1m) and channels (1m). Before transplanting, the recommended doses of fertilizers (N:P:K : 100:60:40) were applied before transplanting of seedlings into main plots. Nitrogen in the form of urea (220kg/ha), Phosphorous in the form of single super phosphate (375kg/ha) and potassium in the form of muriate of potash (66.4kg/ha) were utilized. However, the half of nitrogen was given as top dressing and remaining half of nitrogen was top dressed after 30 days after transplanting and balance half dose was applied at panicle initiation stage. All the recommended agronomic practices were conducted but there was no application of any pesticides during the crop season. Observations were recorded at tillering stage (30DAT), maximum tillering stage (50DAT) and heading stage (90DAT) during the crop season by dissecting fifty infested tillers (deadhearts and white earheads) from each block. The larvae were kept under observation for the confirmation of the species of stem borers of rice. The species were identified based on larval characters described by Kok and Varghese (1966) as mentioned in table- 4.

The same morphological parameters were also utilized by Nishida and Torii (1970), Rai (1984) and Hattori and Siwi (1986) for the identification of stem borer species of rice. After the confirmation of species of stem borers, number of larvae were counted and computed in the form of percentage of each species of the stem borers at respective stages of the crop.

RESULTS AND DISCUSSION

The pooled data presented in Table 3 and Fig. 1. exhibited that all the four species of stem borer of rice viz. yellow stem borer, *Scirpophaga incertulas* (Walker), pink stem borer, *Sesamia inferens* (Walker), white stem borer, *Scirpophaga innotata* (Walker) and dark headed striped borer, *Chilo polychrysus* (Meyrick) were prevalent during *kharif*, 2010 and *kharif*, 2011. Among all the four species, the population of yellow stem borer was found to be dominant over other species of stem borer and recorded 87.00% to 93.00% at 30 DAT during two *kharif* seasons respectively. The population was 82.50% to 89.00% at 50 DAT and 85.00% to 91.50% at 90




Figure 1: Species composition of stem borers of rice (pooled mean of kharif, 2010 and kharif, 2011)

DAT with respect to all the stages (Table 1 and 2) viz. tillering stage, maximum tillering stage and heading stage of the crop growth.

Species composition found during the investigation during *kharif*, 2010 that yellow stem borer recorded higher population 93.00% followed by pink stem borer (3.00%), white stem borer (2.25%) and dark headed striped borer (1.50%) at 30 DAT. Yellow stem borer also showed higher population (89.00%) followed by pink stem borer (5.50%), white stem borer (3.50%) and dark headed striped borer (2.00%) at 50 DAT. At 90 DAT, yellow stem borer also revealed its dominance with higher population (91.50%) followed by pink stem borer (5.00%), white stem borer (2.00%) and dark headed striped borer (1.50%).

It was found during the investigation during *kharif*, 2011 that yellow stem borer recorded higher population (87.00%) followed by pink stem borer (6.00%), white stem borer (4.00%) and dark headed striped borer (3.00%) at 30 DAT. Yellow stem borer also showed higher population (82.50%) followed by pink stem borer (9.00%), white stem borer (5.00%) and dark headed striped borer (3.50%) at 50 DAT. At 90 DAT, yellow stem borer also revealed its dominance with higher population (85.00%) followed by pink stem borer (8.50%), white stem borer (3.50%) and dark headed striped borer (3.00%).

The mean percentage during *kharif*, 2010 and 2011 it was observed that the population of yellow stem borer was dominated with 90.00% followed by pink stem borer with 4.50%, white stem borer with 3.25% and dark headed striped borer with 2.25%. Yellow stem borer also showed higher population (85.75%) followed by pink stem borer (7.25%), white stem borer (4.25%) and dark headed striped borer (2.75%) at 50 DAT. At 90 DAT, yellow stem borer also revealed its dominance with higher population (88.25%) followed by pink stem borer (6.75%), white stem borer (2.75%) and dark headed striped borer (2.25%).

Table 1: Species composition of stem borers of rice during *kharif*, 2010

Stem Borer Species	Percentage Composition (%)			
	30DAT	50DAT	90DAT	Mean
1.Yellow stem borer (<i>Scirpophaga incertulas</i>)	93.00	89.00	91.50	91.16
2.White stem borer (<i>Scirpophaga innotata</i>)	2.50	3.50	2.00	2.67
3.Dark headed striped borer (<i>Chilo polychrysus</i>)	1.50	2.00	1.50	1.67
4.Pink stem borer (<i>Sesamia inferens</i>)	3.00	5.50	5.00	4.50

Table 2: Species composition of stem borers of rice during *kharif*, 2011

Stem Borer Species	Percentage Composition (%)			
	30DAT	50DAT	90DAT	Mean
1. Yellow stem borer (<i>Scirpophaga incertulas</i>)	87.00	82.50	85.00	84.83
2. White stem borer (<i>Scirpophaga innotata</i>)	4.00	5.00	3.50	4.17
3. Dark headed striped borer (<i>Chilo polychrysus</i>)	3.00	3.50	3.00	3.17
4. Pink stem borer (<i>Sesamia inferens</i>)	6.00	9.00	8.50	7.83

Table 3: Species composition of stem borers of rice (pooled mean of *kharif*, 2010 and *kharif*, 2011)

Stem Borer Species	Percentage Composition (%)			
	30DAT	50DAT	90DAT	Mean
1. Yellow stem borer (<i>Scirpophaga incertulas</i>)	90.00	85.75	88.25	88.00
2. White stem borer (<i>Scirpophaga innotata</i>)	3.25	4.25	2.75	3.42
3. Dark headed striped borer (<i>Chilo polychrysus</i>)	2.25	2.75	2.25	2.42
4. Pink stem borer (<i>Sesamia inferens</i>)	4.50	7.25	6.75	6.16

Table 4: The species were identified based on larval characters described by Kok and Varghese (1966)

Species	Head	Body	Prothoracic shield	Crochets of prolegs
1. <i>Scirpophaga incertulas</i> (Yellow stem borer)	Yellowish brown	Creamy yellow 20-25 mm 1 st abdominal segment white	Yellowish brown	Biordinal, sometimes almost uniordinal, arranged in an ellipse.
2. <i>Scirpophaga innotata</i> (White stem borer)		Creamy yellow 20-25 mm	Yellowish brown, anterior margin tinged with dark color	
3. <i>Chilo polychrysus</i> (Dark headed striped borer)	(Dark headed striped borer) Black to blackish brown	Dull white tinged with pink gray with longitu dinal stripes 17-22mm.	Black to blackish brown	Almost triordinal arranged in a circle
4. <i>Sesamia inferens</i> (Pink stem borer)	Reddish brown	Milky white tinged with pink or purple 30-35mm.	Brown	Uniordinal arranged in a longitudinal band

From mean percentage composition of stem borers of rice, it is quite cleared that during *kharif*, 2010 and *kharif*, 2011 yellow stem borer was found dominant with higher population (88.00%) followed by pink stem borer (6.16%), white stem borer (3.42%) and dark headed striped borer (1.42%).

The present findings are in close agreement with the findings of DRR (2008) that yellow stem borer was the dominant species (89.50%) and while pink stem borer, white stem borer and dark headed borer were prevalent during the crop period at Pusa. Similar observations were also reported by Husain and Begum (1985), Damayanti et al. (1991), Rahim et al. (1992), Pathak and Khan (1994), Islam (1996), Catling and Islam (1999), Ragini et al. (2000), JunMing et al. (2003), Lal (2006), Rai et al. (2006), Rahman (2007), DRR (2009), Joshi et al. (2009) and DRR (2011) that yellow stem borer was predominating species throughout the crop season. For preventing the damage rice crop caused by yellow stem borer it is important to control the dominant species to increase agricultural productivity of rice and raise economic condition.

REFERENCES

- Agricultural statistics at a glance, 2011.** On line <http://dacnet.nic.in/eands/latest.html>.
- Ampong-Nayarko., Kwesi and De Dutta, S. K. 1991.** *A hand book for weed control in rice.* IRRI, Philippines. p.7.
- Atwal, A. S. and Dhaliwal, G. S. 2005.** *Agricultural Pests of South Asia and their Management.* Kalyani Publishers, New Delhi. pp.181-182.
- Balakrishna, B. and Satyanarayana, P. V. 2013.** Genetics of brown

planthopper (*Nilaparvata lugens* Stal.) Resistance in elite donors of rice (*Oryza sativa* L.). *The Bioscan.* **8(4):** 1413-1416.

Catling, H. D. and Islam, Z. 1999. Pests of deep water rice and their management. *Integrated Pest Management Reviews.* **4:** 193-229.

Cramer, H. H. 1967. Plant protection and world cup protection: Pflanzenschutz. *Nachar.* **20(1):** 524.

Damayanti, D., Siwin, S. S. and Soejitno, J. 1991. Distribution and composition of rice stem borer species in central Java. Balai Penelitian Tanaman Pangan Bogor (Indonesia). *Seminar Hasil Penelitian Tanaman Pangan Balittan Bogor Tahun.* **2:** 378-387.

DRR, 2009. *DRR Annual Progress Report, 2009,* Directorate of Rice Research, Rajendranagar, Hyderabad. **2:** 2.19-2.35, 2.50-2.56, 2.57-2.60.

DRR, 2011. *DRR Annual Progress Report, 2011,* Directorate of Rice Research, Rajendranagar, Hyderabad. **2:** 2.21-2.35, 2.52-2.65.

Gupta, S. P., Pandey, U. S., Pandey, V. and Mishra, C. H. 2002. Screening of rice cultivars under wet condition against major insect pests of rice. *Shashpa.* **9(2):** 189-190.

Hattori, I. and Siwi, S. 1986. Rice Stem Borers in Indonesia. *Japan Agricultural Research Quarterly.* **2(1):** 25-30.

Husain, M. and Begum, N. 1985. Seasonal stem borer (SB) population fluctuations in Mymensingh, Bangladesh. *International Rice Research Newsletter.* **10:** 22.

Islam, Z. 1996. Yellow stem borer, a threat to boro rice in coastal belt of Bangladesh. *J. Entomology.* **6(1&2):** 45-52.

Joshi, G., Ram, L. and Singh, R. 2009. Population dynamics of paddy stem borers in relation to biotic and abiotic factors. *Annals of Biology.* **25(1):** 47-51.

JunMing, S., Gang, W., XianWen, Z., Qian, X., ShanZhang, L. and ChengFa, S. 2003. The population dynamics of the yellow rice borer causes of outbreaks and control strategy. *Entomological Knowledge.*

40(2): 124-127.

Kok, L.T. and Varghese, C. 1966. The four major lepidopterous rice stem borers in Malaya. *Malaya Agriculture J.* **45(3):** 275-288.

Lal, R. 2006. Effect of sowing method of wheat after rice on the hibernating larvae of rice stem borers vis-à-vis their management in Horgana. *Indian J. Agrometeorology.* **68(3):** 225-229.

Manikandan, N., John Samuel Kennedy and Geethalakshmi, V. 2014. Life History And Population Dynamics Of Rice Leaf folder at Different Temperatures. *The Ecoscan.* **8(3&4):** 315-320.

Nishida, T. and Torii. 1970. Identification of rice stem borers. A hand book of field methods for research on rice stem borers and their natural enemies. *IBP Hand Book.* **(14):** 7-41.

Pathak, M. D. 1975. *Insect Pests of Rice.* International Rice Research Institute, Los Banos, Philippines. pp.68.

Pathak, M. D. and Dhaliwal, G. S. 1981. *Trends and strategies for rice insect problems in Tropical Agriculture, IRRI, Research Paper series No. 64.* pp.15.

Pathak, M. D. and Khan, Z. R. 1994. *Insect Pests of Rice.* International Rice Research Institute Publication, Philippines. p. 89.

Ragini, J. C., Thangaraju, D. and David, P.P. M. 2000. Stem borer species composition in Tamil Nadu, India. *International Rice Research Newsletter.* **25: 15.**

Rahim, A., Jabbar, A., Hashmi, A.A. and Khan, N.A. 1992. Chemical control of yellow stem borer in Sindh. *Pakistan J.Agricultural Research.* **13(4): 380-389.**

Rahman, M. M. 2007. The effect of variety, age and climate on the abundance of rice insect pests and their natural enemies. *J. Subtropical Agricultural Research and Development.* **5(1/2): 239-245.**

Rai, A. K. 1984. Bionomics and losses due to stem borers in deep water rice. *M.Sc. Thesis.* Department of Entomology, RAU, Pusa.

Rai, C. P., Mishra, A. K. and Choudhary, N. K. 2006. Incidence of rice stem borer species in varying ecological amplitudes of Bihar. *J. Applied Biology.* **16 (1/2): 53-56.**

Rao, V. P. 1965. Natural enemies of rice stem borer and allied species in various parts of the world and possibilities of their use in biological control of rice stem borers in Asia. *Technical bulletin commonwealth institute of biological control.* **6: 1-68.**

Roy, B., Surje, D. T. and Mahato, S. 2013. Biodiversity of farmers' varieties of rice (*Oryza sativa* L.) at repository of uttarbanga krishi viswavidyalaya: A reservoir of important characters. *The Ecoscan. Special Issue.* **4: 145-151.**

Saxena, S. and Murty, N. S. 2014. Weather based model development for outbreak of mustard aphid (*I. erysimi.,kalt*) using artificial neuralnetwork. *The Ecoscan.* **8(1&2): 47-52.**