

# USE OF HYPERSPECTRAL REMOTE SENSING FOR QUANTITATIVE ESTIMATION OF SOIL ORGANIC CARBON

SHWETA GAUTAM<sup>1\*</sup>, A. S. NAIN<sup>1</sup> AND POONAM GAUTAM<sup>2</sup>, HIMANI BISHT<sup>1</sup>

<sup>1</sup>Department of Agrometeorology,

<sup>2</sup>Department of Soil Science,

College of Agriculture, GBPUAT, Pantnagar 263 145 ,U.S. Nagar, Uttarakhand, INDIA.

e-mail: gautamshweta808@gmail.com

## INTRODUCTION

During last decades there has been a rapid extension in technologies for agriculture advancement including Global Positioning System (GPS), Geographic Information Systems (GIS), remote sensing and crop simulation models (Garofalo *et al.*, 2009; Hbirkou *et al.*, 2012 and Puranik *et al.*, 2014). Organic carbon (OC) plays a major role in keeping fertility of soil and thereby flourishing the biological activity (Gebbers *et al.*, 2010 and Rajan 2011). Conventional techniques for identifying soil organic carbon (SOC) require very dense sampling and need more time for characterization of spatial variability within an area. However imaging spectroscopy technology in the visible (VIS) and near-infrared (NIR) region is a more efficient, faster and less expensive that provides data with high temporal and spatial resolution (Simbahan *et al.*, 2006, Stevens *et al.*, 2010). Hyper spectral images due to high number of spectral bands and improved spatial and radiometric resolution provides more information than traditional multispectral images. Several studies have used hyperspectral remote sensing imageries to estimate soil properties (Ray *et al.*, 2004; Gomez *et al.*, 2008; and Nowkandeh *et al.* 2013). The present hyperspectral sensors can be used to map the characteristic of soil properties with great accuracy. Therefore in the present study, an attempt has been made for mapping of soil organic carbon using hyperspectral remote sensing in Mollisols of Uttarakhand.

## MATERIALS AND METHODS

### Study area

The study was carried at Agricultural farm GBPUA&T, Pantnagar, U.S. Nagar (Uttarakhand) which is located at 29°N latitude, 79.29°E longitude and with an altitude of 243.80m from the mean sea level in the *Tarai* belt, about 30 km southward of foot hills of the Shivalik range of Himalayas. Hyperion image covering whole University Farm including Nagla farm, Vegetable Research Centre, Crop Research Centre and Horticulture Research Centre etc has been shown in figure 1. The study area falls under sub-humid to sub-tropical climate with hot dry summers and cool winters.

### Data Acquisition

#### Remote Sensing Data Acquisition

The Cloud free Hyperion imagery of May 2<sup>nd</sup>, 2013 was acquired from the United States Geological Survey (USGS) archive (<http://earthexplorer.usgs.gov/>). Hyperion, the first spaceborne Hyperspectral sensor was launched on 21<sup>st</sup> November 2000 as part of NASA's New Millennium Program. Hyperion has 242 spectral bands spanning a spectral range from 0.4 to 2.5  $\mu\text{m}$ , with a sampling interval of 10nm. The imagery was received as a full long scene (185-km strip) and at level 1 (L1GST) processing level in GeoTIFF format, written as band-interleaved-by-line

## ABSTRACT

Soil organic carbon (SOC) appears as a promising indicator to know the overall fertility status of soil. In the present investigation, a remote sensing based approach for estimation and mapping of soil organic carbon using hyperspectral remote sensing imagery was used. Soil samples (Mollisols) were collected from the agricultural farm of GBPUA & T, Pantnagar, Uttarakhand, India. VIS-NIR spectra were collected over the sampling points. Using hyperion sensor VIS-NIR spectra was transformed into different spectral indices like Brightness Index (BI), Hue Index (HI), Saturation Index (SI), Coloration Index (CI), Redness index (RI) Normalized Difference Vegetative Index (NDVI) and Ratio Vegetation Index (RVI). Results indicated that soil organic carbon was significantly correlated with spectral indices *viz* blue (487.86nm), green (569.27nm), SWIR (2345.1nm) bands and brightness index. Multiple regression models were also developed to estimate soil organic carbon based on spectral indices. The comparison between observed and estimated SOC showed quite good agreement with  $R^2$  of 0.83 in multivariate model-10 and RMSE of 0.15. Later, multivariate model was used to generate the layer of SOC representing spatial distribution. This showed the capability of hyperspectral remote sensing for estimation of SOC which can act as a base map to predict management practices in precision agriculture.

## KEY WORDS

Hyperion  
Hyperspectral remote sensing  
Soil organic carbon

Received : 14.05.2016

Revised : 10.07.2016

Accepted : 13.08.2016

\*Corresponding author

(BIL) files stored in 16-bit signed integer radiance values. The L1GST product is radiometrically corrected, geometrically resampled, and registered to a geographic map projection with elevation correction applied.

**Soil Data**

Soil samples were collected from 25 locations of university farm using hand held khurpi. To get good representation of site, four soil samples were collected from a field at 0-15cm depth. Thereafter, all four soil samples were properly mixed and half a kilo soil samples has been taken for further processing. The field work was undertaken during the month of May on a clear sunny day. The soil organic carbon content was determined by Walkley and Black (1934) method.

**Pre-processing and Atmospheric Correction of hyperion image**

The Hyperion EO-1 sensor records radiance of 242 spectral bands with band width of 10nm. In order to standardize L1GST hyperion image into ENVI format band, the original image was imported using Hyperion Tools 2.0 (White 2013) that contains wavelength, full width half maximum and bad band information.. Although Hyperion Tools 2.0 also flags bad bands, visual inspection revealed additional bands dominated by noise. In this study, 147 bands were used finally used which are: bands 9 – 55, 86 – 119, 133 – 164, 183 – 184, 188 – 200, and 202 – 220. The image was atmospherically corrected into the surface reflectance using QUAC (Quick atmospheric correction technique).

**Computation of Spectral Indices**

Various soil-related spectral indices were computed from Hyperion data, after converting the digital numbers into radiance values. Those indices included soil related indices such as, Brightness Index (BI), Hue Index (HI), Saturation Index (SI), Coloration Index (CI), Redness index (RI) Normalized Difference Vegetative Index (NDVI) and Ratio Vegetation Index (RVI). The statistical models for estimating these indices are presented in Table 1.

Where R = 681.19nm, G = 569.27nm, B = 487.87nm, NIR = 894.88nm

**Development of Multivariate model**

In the present study, stepwise multivariate statistical regression was carried out using SPSS package to model the relationship between spectral indices and soil organic carbon. The representative expression of multivariate model has been mentioned below:

$$Y = b_0 + b_1X_1 + b_2X_2 + \dots + b_nX_n$$

Where, Y = dependent variables, b0 = estimated constant, bn = estimate coefficients, Xn = independent variables

Spectral indices mentioned in Table 1 together with different individual's bands were used as independent variables, while SOC has taken as dependent variable.

**Statistical Analysis**

**Correlation Coefficient**

A relationship was established between observed and estimated SOC to analyze the accuracy of multivariate model.

$$R = \frac{\sum_{n=1}^N (R_n - R') (C_n - C')}{\sqrt{\sum_{n=1}^N (R_n - R')^2 * (C_n - C')^2}}$$

Where, r is the correlation coefficient, R is the selected variables (spectral band values and indices, N is the number of soil samples, here N is 25, Cn is soil organic carbon content of sample n and R as well as C' are the mean values.

**Root Mean Square Error (RMSE)**

The RMSE has been used as a criterion for model evaluations and has been computed as

$$RMSE = \sqrt{\frac{\sum_{i=1}^n (X_{obs,i} - X_{model,i})^2}{n}}$$

where Xobs is observed values and Xmodel is modelled values

**Estimation of Soil Organic Carbon**

Model was developed by analyzing the relationship between SOC and spectral indices derived using Hyperion hyperspectral reflectance image and was finally used for mapping the distribution of SOC. The complete approach of estimating spatial distribution of SOC has been presented in figure 2.

**RESULTS AND DISCUSSION**

**Laboratory analysis results**

Results of chemical analysis of 25 soil samples showed high spatial variability in SOC content among the different soil sampling sites. This may be due to natural or by human activity (Olorunlana, 2015). The soils of Pantnagar belong to Mollisols order according to U.S.D.A. Taxonomic System (Soil Survey Staff, 1978) and have six soil series viz. Patharchatta sandy loam, Nagla loam, Haldi loam, Phoolbagh clay loam, Khamia




Figure 1. Location of study area and Hyperion image of 2<sup>nd</sup> May, 2013 showing agriculture farm of the GBPUA&T, Pantnagar


**Table 1: Spectral indices calculated using hyperion image for soil organic carbon study**

Index	Formula Index	References
BI	$\{(B^2 + G^2 + R^2)/3\}^{1/2}$	(Mathieu and Pouget, 1998)
RI	$R_2/(B * G_3)$	(Mathieu and Pouget, 1998)
SI	$(R-B)/(R+B)$	(Mathieu and Pouget, 1998)
CI	$(R-G)/(R+G)$	(Mathieu and Pouget, 1998)
HI	$(2 * R - G - B)/(G - B)$	(Mathieu and Pouget, 1998)
RVI	NIR/R	(Jordan, 1969)
NDVI	$(NIR - R)/(NIR + R)$	(Rouse et al., 1974)

CDI (Carbon difference index) was developed by taking the difference between band199 and band184  
 Where, R, G, B and NIR are red, green, blue and near infrared bands respectively

**Table 2: Laboratory analysis of soil properties**


Parameters	Minimum	Maximum	Mean	Median	Std. Dev.	C.V. (%)
SOC (%)	0.37	1.77	.79	.63	0.37	47



**Figure 2: A flow chart summarizing the analytical and technical stages**

**Table 3: Correlation study of spectral parameters derived from Hyperion image and SOC**

S. No.	Spectral Parameters	Coorelation with SOC
1	Blue (487.86nm)	-0.375
2	Green (569.27nm)	-0.346
3	Red (681.19nm)	-0.287
4	Band 97 (1114.2nm)	-0.08
5	Band 114 (1578.3nm)	-0.382
6	Band 201(2163.4nm)	-0.382
7	Band215 (2304.7nm)	-0.384
8	Band 219 (2345.1nm)	-0.503
9	RVI	-0.1
10	NDVI	-0.033
11	CDI	0.05
12	HI	0.037
13	BI	-0.420
14	CI	0.045
15	RI	-0.043
16	SI	-0.041



**Figure 3: A scatter plot presenting the correlation against predicted SOC and measured SOC (n= 25)**

sandy loam and Beni silty clay loam (Deshpandey et al., 1971). The statistical summary (Table 2) of above soil samples of different locations showed the minimum, maximum, mean, median, standard deviation and coefficient of variation (CV).

SOC content of samples ranged between 0.37% to 1.77% with mean value of 0.98%. Variability of soil was analyzed by estimating coefficient of variation (CV) and it was found 47% with the standard deviation (Std. Dev.) of 0.37.

**Pearson correlation and Multiple regression test results**


The statistical analysis was performed using SPSS package between soil organic carbon and different spectral indices (Table 3). As SOC increases, the soil appears darker and consequently, the reflectance from soil surface will decrease and hence in general there is a negative correlation between SOC and reflectance of Hyperion bands. Furthermore, as wavelength increases from visible to Short wave infrared (SWIR) the correlation shows an ascending (-0.08 in band 97) and then a descending (-0.38 in band 114, -0.38 in band 215, -0.50 in band 219) pattern. Similar findings were reported by Ladoni, et al. (2010) and Nowkandeh et al. (2013). The highly significant correlations with blue band (-0.37), green bands (-0.35), brightness index (-0.42) and SWIR in SOC showed conformity with study conducted by Chen et al. (2000) and Ray et al. (2004).

**Multivariate statistical regression models for organic carbon SOC**

The multivariate models were developed between SOC and spectral parameters using stepwise regression technique. Ten

**Table 4: Empirical equations between organic carbon and spectral parameters derived using stepwise regression technique**

S. No.	Model	R <sup>2</sup>
1	O.C. (%) = 1.18-1.46* Band 219	0.25
2	O.C. (%) = 1.79-1.695*Band219-1.593*Band97	0.31
3	O.C. (%) = 2.03-2.35*Band219-3.53*Band97+2.32*Band33	0.35
4	OC (%) = 2.302-2.17*Band219-3.48*Band97-3.82*Band33-3.10*BI	0.39
5	O.C. (%) = 3.36-1.34*Band219-3.34*Band97+2.94*Band33-9.15*BI+61.59*RI	0.53
6	O.C. (%) = 4.53-1.34*Band219-6.11*Band97+2.72*Band33-12.66*BI+119.26*RI+0.27*RV1	0.65
7	O.C. (%) = 4.65-2.06*Band219-5.68*Band97+1.64*Band33-15.73*BI+141.54*RI+0.39*RV1+2.14*Band215	0.71
8	O.C. (%) = 4.69-1.69*Band219-4.78*Band97+1.64*Band33-17.32*BI+150.82*RI+0.32*RV1+2.07*Band215+0.13*CDI	0.73
9	O.C. (%) = 5.0-1.52*Band219-5.22*Band97+2.15*Band33-17.63*BI+150.43*RI+0.27*RV1+1.64*Band215+0.18*CDI+.001*HI	0.77
10	O.C. (%) = 4.55-1.36*Band219-2.20*Band97+0.29*Band33-20.07*BI+162.46*RI+0.68*RV1+1.86*Band215+0.24*CDI+.002*HI-2.35*NDVI	0.83

**Figure 4: Spatial distribution of SOC**

models were developed for SOC and providing R<sup>2</sup> value from 0.25 for model-1 to 0.83 for model-10. The reasonably good multiple regression model were noticed from model-6 (R<sup>2</sup>=0.65) to model-10 (R<sup>2</sup>=0.83). Model-1 (R<sup>2</sup>=0.25), model-2 (R<sup>2</sup>=0.31), model-3 (R<sup>2</sup>=0.35) model-4 (R<sup>2</sup>= 0.39) and model-5 (R<sup>2</sup>= .53) did not exhibit significant results. The most appropriate multivariate model for SOC estimation was model-10 having coefficient of determination (R<sup>2</sup>) as 0.83 and RMSE has been observed as .15. Srivastava, *et al.*, 2004; Kadupitiya *et al.*, 2010; Sahoo, *et al.*, 2012; Ghosh *et al.*, 2012 obtained good calibration for SOC based on spectral reflectance.

A scatter plot exhibiting one to one relationship between observed and predicted SOC (from model-10) values has been shown figure 3. Results showed a good agreement between the predicted values and the measured values. Ray *et al.*, 2004 and Luo *et al.*, 2008 calculated several soil color indices namely brightness index, coloration index, hue index, redness index and saturation index using Hyperion Hyperspectral data for statistical modeling of SOC. An overall accuracy of 83% was achieved showing the significance of the technique.

#### Variability map generation

Based on the statistical analysis, multivariate model was used to map the spatial distribution of SOC. SOC variability map was generated using hyperion hyperspectral image with the help of the model-10 having R<sup>2</sup> value of 0.83. The band math function embedded in ENVI 4.8 software has been used to generate SOC layer (Fig.4). The values of SOC content vary from 0.37 to 1.77% (Table 2) over the agriculture farm region of Pantnagar. Most of the regions come under the range of

high organic carbon content (*i.e.* 0.7-1.3%, Deshpande *et al.*, 1971). The spatial distribution of SOC can be used for site specific soil management. In the past also (Zheng, 2008, Wang *et al.*, 2010) remote sensing techniques have been used to generate soil fertility map, which also exhibited similar kind of SOC distribution. In order to achieve higher accuracy it would be appropriate to collect soil samples at the time of satellite pass. Further, research is recommended for different locations, climate, vegetation soil types and surface conditions.

#### REFERENCES

- Deshpandey, S. B., Fehrenbacher, J. B. and Ray, B. W. 1971. Mollisol of Tarai region of Uttar Pradesh, Northern India, *Genesis and Classification*, Geoderma. **6**: 195-201.
- Garofalo, P., Di Paolo, E. and Rinaldi, M. 2009. Durum wheat (*Triticum durum* Desf.) in rotation with faba bean (*Vicia faba* var. minor L.): long-term simulation case study. *Crop and Pasture Science*. **60**: 240-250.
- Gebbers, R. and Adamchuk, V. I. 2010. *Precision Agriculture and Food Security*. Science. **327**: 828-831.
- Ghosh, R., Padmanabhan, N., Patel, K. C. and Siyolkar, R. 2012. Soil fertility parameter retrieval and mapping using hyperion data. In *Investigations on Hyperspectral Remote Sensing Applications* (eds Panigrahy, S. and Manjunath, K. R.), Space Applications Centre (ISRO), Ahmedabad, pp.29-31.
- Hbirkou, C., Patzold, S., Mahlein, A.K. and Welp, G. 2012. Airborne hyperspectral imaging of spatial soil organic carbon heterogeneity at the field-scale. *Geoderma*. pp.175-176. 21-28.
- Jordan, C. F. 1969. Derivation of leaf area index from quality of light on the forest floor. *Ecology*. **50**: 663-666.
- Kadupitiya, H. K., Sahoo, R. N., Ray, S. S., Chakraborty, D. and Ahmed, N. 2010. Quantitative assessment of soil chemical properties using visible (VIS) and near-infrared (NIR) proximal hyperspectral data. *Trop. Agric*. **158**: 41-60.
- Ladoni, M., Alavaipah, S. K., Bahrami, H. A. and Noroozi, A. A. 2010. Remote sensing of soil organic carbon in semiarid region of Iran. *Arid Land Research and Management*. **24**: 271-281.
- Mathieu, R. and Pouget, M. 1998. Relationship between satellite-based radiometric indices simulated using laboratory reflectance data and typical soil colour of an arid environment. *Remote Sensing of Environment*. **66**:17-28.
- Nowkandeh, S. M., Homaee, M. and Noroozi, A. A. 2013. Mapping Soil Organic Matter Using Hyperion Images. *International J. Agronomy and Plant Production*. **4(8)**:1753-1759.
- Olorunlana, F. A. 2015. Factor analysis of soil spatial variability in Akoko Region of Ondo State. *J. Geography and Regional Planning*. **8(1)**: 12-15.
- Puranik, H. V., Nain, A. S. and Murty, N. S. 2014. GIS based suitability

evaluation of land for acacia catechu in degraded and waste lands in Uttarakhand. *The Ecoscan*. **8(3and4)**: 321-325.

**Rajan, D. S. and Bindhu, R. 2011.** Distribution pattern of organic carbon in the sediments of ashtamudi estuary. *The Ecoscan, Special issue.1*: 71-74.

**Ray, S. S., Singh, J. P. Das, G. and Panigrahy, S. 2004.** Use of High Resolution Remote Sensing Data for Generating Site-specific Soil Mangement Plan. XX ISPRS Congress, Commission 7. Istanbul, Turkey The International Archives of the Photogrammetry, *Remote Sensing and Spatial Information Sciences*. pp.127-131.

**Rouse, J. W., Haas, R. H., Schell, J. A., Deering, D. W. and Harlan, J. C. 1974.** Monitoring the Vernal Advancements and Retroradation (Green wave Effect) of Nature Vegetation. NASA/GSFC Final Report, NASA, Greenbelt, MD, pp.371.

**Sahoo, R. N., Ray, S. S., Chopra, U. K. and Govil, V. 2012.** Estimation of soil parameters using ground and space based hyperspectral data. In Investigations on Hyperspectral Remote Sensing Applications (eds Panigrahy, S. and Manjunath, K. R.), Space Applications Centre (ISRO), Ahmedabad, pp. 43-51.

**Simbahan, G. C., Dobermann, A., Goovaerts, P., Ping, J. L. and Haddix, M. L. 2006.** Fine resolution mapping of soil organic carbon based on multivariate secondary data. *Geoderma*. **132**: 471-489.

**Soil Survey Staff. 1978.** Soil Texonomy. U.S.D.A. Handbook No.436.

**Srivastava, R., Prasad, J. and Saxena, R. 2004.** Spectral reflectance properties of some shrink-swell soils of Central India as influenced by soil properties. *Agropedology*, 2004, 14,pp. 45-54.

**Stevens, A., Udelhoven, T., Denis, A., Tychon, B., Lioy, R., Hoffmann, L., Wesemael, B. and Adkins, W. 2010.** Measuring soil organic carbon in croplands at regional scale using airborne imaging spectroscopy. *Geoderma*. **158**: 32-45.

**Walkley, A. and Black, I. A. 1934.** An examination of the Degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science*. **37**: 29-38.

**Wang, J., He, T., Lv, C., Chen, Y. and Jian, W. 2010.** Mapping soil organic matter based on land degradation spectral response units using Hyperion images. *International J. Applied Earth Observation and Geoinformation*. **12**: S171-S180.

**White, D. 2013.** Hyperion Tools 2.1. [online]. Available from: <http://www.exelisvis.com/UserCommunity/CodeLibrary.aspx>

**Zheng B, 2008.** Using satellite hyperspectral imagery to map soil organic matter, total nitrogen and total phosphorus. MSc thesis of Indiana University, Department of earth science.

