

MULTIVARIATE ANALYSIS FOR TRAIT ALLIANCE OF BREAD WHEAT YIELD UNDER TERMINAL HEAT STRESS CONDITIONS

SWARUPA NANDA MANDAL^{1,2}, SATYAVIR SINGH DHANDA¹, RENU MUNJAL^{1*} AND CHANCHAL PRAMANIK³

¹Department of Genetics & Plant Breeding, CCS Haryana Agricultural University, Hisar - 125 004, INDIA

²Department of Genetics & Plant Breeding,

Extended Campus of Bidhan Chandra Krishi Viswavidyalaya, West Bengal - 713 101, INDIA

³Tata Consultancy Services, Hyderabad - 500 034, INDIA

e-mail: munjalrenu66@gmail.com

INTRODUCTION

Heat stress during post-anthesis growth stages predominantly influences assimilates availability, translocation of photosynthates to the grain and starch synthesis and deposition in the developing grain (Modarresi *et al.*2010). The net result is a lower grain yield because of lower grain weight (Gibson and Paulsen, 1999; Pandey *et al.* 2015). It is difficult to make progress for grain yield and yield components under heat stress conditions as these are highly intricate in nature and entails several tolerance and avoidance mechanisms. A detailed understanding of the genetics and physiology of grain yield and its components under heat stress conditions would improve the efficiency of breeding programs by identifying relevant indices for selecting varieties (Kalyar *et al.*2014).

Among the heat stress related traits, cell membrane is a major physiological parameter which disrupts due to heat stress thus the lipid bilayer membranes become more fluid by either due to denaturation of proteins or an intensification of unsaturated fatty acids (Savchenko *et al.*2002) and measurement of solute leakage from plasma membrane or membrane thermo stability can be used to estimate the heat stress tolerance (Blum, 2001; Dhanda and Munjal, 2006). Canopy temperature depression (CTD) is another trait involved in a number of different physiological and metabolic processes including stomatal conductance, photosynthetic rate, vascular capacity, transpiration rate, stomatal conductance and may be indicative of heat avoidance capability of plant through keeping the canopy cooler during high demand for transpiration, and photo assimilation caused by rapidly filling kernels (i.e. sink strength) and good vascular system. Hence, CTD is positively associated with yield under heat stress (Bahar *et al.* 2011).

Statistical techniques have important role in establishing relationship between the yield and its related components. Though correlation coefficient is an important statistical method in selection for higher yield (Steel and Torrie, 1960), may not provide adequate details about the relationship among different variables (Del Moral *et al.*2003), as much as multivariate statistical methods give. Multiple linear regression, stepwise regression, path analysis, principal component analysis and cluster analysis have been established as useful methods to decipher these relations in crop plants (Saed-Moucheshi *et al.*2013). The multivariate statistical analyses can furnish insights the structure of data and traits' alliance. This study was directed to describe and interpreting the association between wheat grain yield and its related components under terminal heat stress conditions with aiming to provide theoretical foundations guiding wheat breeders for researching the alliance of the main yield components and their influences on wheat plant productivity. Thus the present investigation was initiated to examine the influence of high temperature stress at anthesis and post anthesis stages of plant growth, to

ABSTRACT

One hundred twenty five bread wheat recombinant inbred lines were subjected to late heat stress under field conditions during the year 2010-11. The experiment was conducted at CCS Haryana Agricultural University Hisar using randomized complete block design with three replications under late sown conditions. The correlation coefficients revealed that the genotypes having high grain yield also had more number of yield components, namely, number of tillers per plant, number of grains per plant, grain weight per spike, number of grains per spike. These genotypes also had high biomass, harvest index and better in terms of heat stress related parameters, i.e., canopy temperature depression and membrane thermo stability and earlier in heading and maturity. Path coefficient analysis indicated that the grain weight/spike had the highest direct contribution towards grain yield per plant followed by number of grains/plant, number of tillers/plant, biomass and harvest index. Multiple regression analyses revealed that number of tillers per plant was the most contributing trait followed by numbers of grains per plant, grain weight per spike, number of grains per spike, biomass per plant and harvest index. The diversity analysis indicated that the genotypes in one cluster were in general different from another in terms of their response to heat stress.

KEY WORDS

Canopy temperature depression
Cluster analysis
Membrane thermo stability
Multiple linear regression

Received : 11.10.2015

Revised : 17.01.2016

Accepted : 19.02.2016

*Corresponding author

identify important traits contributing heat tolerance and promising combination of traits which may be used as the source of heat tolerance in wheat improvement programme.

MATERIALS AND METHODS

Creation of heat stress environments

The experiment was laid out in randomized complete block design with three replications at CCS HAU, Hisar (29°10'N lat., 75°46'E long., 215 m alt.) during 2010-2011. The plot size was of single row of 2.5 m in length with the spacing of 10 cm × 22.5 cm. In order to create heat stress during anthesis and reproductive stages, the sowing was delayed by about one and half month from normal period of sowing, i.e., in last week of December. Mean weekly maximum and minimum temperatures over the growth period were recorded from Agro-Meteorological Observatory, CCS Haryana Agricultural University, Hisar, India (Fig. 1).

Plant materials

The present investigation was carried out on 125 RILs of bread wheat derived from the cross WH 730 (a thermo-tolerant variety) and WH147 (a thermo-sensitive variety). Observations were recorded as the average of five competitive plants selected randomly from each.

Canopy temperature depression

A hand held infrared thermometer, (model AG-42, Tele temp crop, Fullerton CA) used for instantaneous measurement of canopy minus air temperature as canopy temperature depression. Observations were recorded between 12:00 hrs. to 14:00 hrs.

Membrane thermo stability

For membrane thermo stability, method given by Sullivan (1972), modified later by Ibrahim and Quick (2001) was followed. Membrane thermo stability was measured by the formula given below.

$$MTS = 1 - \frac{T_1}{T_2} \times 100$$

T₁ = initial conductance value taken after incubation of leaf discs in a controlled temperature water bath at 49°C for 45

min followed by cooling.

T₂ = final conductance value taken after autoclaved at 0.01 M Pa pressure for 10 min to release all the electrolytes from the leaf discs

Statistical Analysis

Correlation coefficients The phenotypic coefficients of correlation between two characters were determined by using the variance and covariance components as suggested by Al-Jibouri *et al.* (1958).

Multiple linear regressions

Multiple linear regression and partial coefficient of determination (R²) was estimated for each yield component (Snedecor and Cochran, 1981) in order to assess the relative contribution and to develop the prediction model for grain yield.

Stepwise regression

Stepwise regression was used in order to determine the variable accounting for the majority of total yield variability as suggested by Draper and Smith, 1981.

Principal components analysis

Principal component analysis (PCA) was used to reduce data dimensionality by covariance analysis between variables (Everitt and Dunn, 1992).

Cluster analysis

This analysis was performed using a measure of similarity levels and Euclidean distance (Everitt, 1993).

Path coefficient analysis

Path coefficient analysis was carried out using phenotypic and genotypic correlation values of yield components as suggested by Dewey and Lu (1959).

All statistical analyses were performed using SAS (SAS Institute, 2009) and statistical language R (R Core Team, 2013).

RESULTS

Phenotypic correlation analysis

Results revealed that grain yield per plant exhibited positive significant correlations with number of tillers per plant (0.58**), number of grains per plant (0.78**), grain weight per spike

Table 1: Phenotypic correlation coefficients between various characters in recombinant inbred lines of bread wheat under heat stress conditions

Variables	Grain yield /plant	No. of tillers/ plant	No. of grains/ plant	Grain weight/ spike	No. of grains/ spike	Biomass	Plant height	HI	DTH	DTM	CTD
No. of tillers/plant (X1)	0.58**	1									
No. of grains/plant (X2)	0.78**	0.52**	1								
Grain weight/spike (X3)	0.43**	-0.40**	0.31**	1							
No. of grain/spike (X4)	0.33**	-0.29**	0.58**	0.79**	1						
Biomass (X5)	0.82**	0.65**	0.66**	0.20*	0.15	1					
Plant height (X6)	0.41**	0.25**	0.28**	0.16	0.09	0.41**	1				
HI (X7)	0.48**	0.03	0.37**	0.48**	0.38**	-0.05	0.08	1			
DTH (X8)	-0.23**	-0.07	-0.19*	-0.20*	-0.18*	-0.16	-0.35**	-0.14	1		
DTM (X9)	-0.23**	-0.07	-0.20*	-0.21*	-0.19*	-0.16	-0.33**	-0.14	0.95**	1	
CTD (X10)	0.20*	0.17	0.24**	0.06	0.13	0.19*	-0.05	0.08	0.09	0.05	1
MTS (X11)	0.18*	0.16	0.19*	0.02	0.05	0.13	< 0.01	0.1	-0.18*	-0.20*	0.22*

*, **: Significant at 5% and 1% level of probability, respectively

Table 2: The regression coefficient (b), standard error (SE), t-value and probability of the estimated variables in predicting wheat grain yield by the multiple linear regression analysis

Variables	DF	Parameter estimate	Standard error	t-value	Pr > t
X1	1	0.06425	0.06907	0.93	0.3542
X2	1	0.00537	0.00111	4.85	<.0001**
X3	1	1.21177	0.32507	3.73	0.0003**
X4	1	-0.03024	0.00621	-4.87	<.0001**
X5	1	0.21503	0.01799	11.95	<.0001**
X6	1	0.00277	0.00332	0.83	0.4062
X7	1	21.89501	1.73669	12.61	<.0001**
X8	1	0.03633	0.03313	1.10	0.2751
X9	1	-0.04531	0.03304	-1.37	0.1729
X10	1	-0.01486	0.02932	-0.51	0.6134
X11	1	0.00113	0.00449	0.25	0.8019

*, **: Significant at 5% and 1% level of probability, respectively. Y-intercept (a) = 4.53647; SE = 1.52, R² = 0.9806, Root MSE = 0.36393, Adj R² = 0.9788.

Table 3: Relative contribution (partial and model R²) in predicting wheat grain yield, F-value and probability by the stepwise procedure analysis

Step	Variable entered	Partial R ²	Model R ²	SE of estimates
1	X5	0.6963	0.6963	0.00481
2	X7	0.2791	0.9754	0.75078
3	X9	0.0005	0.9758	0.00891

Table 4: Regression coefficient (b), standard error (SE), F-value and probability (significance) of the accepted variables that can be used to predict wheat grain yield by the stepwise procedure

Variables	Coefficient of regression (B)	Standard error (SE)	F value	Significance
X5	0.28057	0.00481	3397.64	<.0001**
X7	27.70354	0.75078	1361.59	<.0001**
X9	-0.01382	0.00891	2.40	0.1236

*, **: Significant at 5% and 1% level of probability, respectively

Table 5: Eigen value of the correlation matrix for the estimated variables of wheat using the principal component procedure

Variables	PC1	PC2	PC3	PC4	PC5	PC6	PC7	PC8	PC9	PC10	PC11
X1	0.202	0.576	0.076	0.039	0.274	-0.151	-0.047	0.061	0.650	0.043	0.304
X2	0.430	0.152	0.300	0.099	0.114	-0.086	-0.357	-0.373	-0.529	-0.042	0.351
X3	0.295	-0.472	0.153	0.137	-0.144	0.174	0.083	0.523	0.100	-0.003	0.553
X4	0.323	-0.398	0.266	0.088	-0.190	0.036	-0.350	-0.339	0.486	0.044	-0.382
X5	0.351	0.392	0.107	0.217	-0.205	0.119	-0.130	0.554	-0.196	-0.012	-0.496
X6	0.299	0.131	-0.210	0.410	-0.168	0.390	0.592	-0.387	0.025	-0.014	0.016
X7	0.268	-0.244	0.228	-0.037	0.721	-0.172	0.409	0.085	-0.077	-0.005	-0.292
X8	-0.355	0.079	0.525	0.167	0.030	0.235	0.054	-0.019	0.064	-0.707	-0.002
X9	-0.356	0.078	0.512	0.204	0.045	0.252	0.055	-0.017	-0.040	0.702	0.016
X10	0.124	0.111	0.404	-0.419	-0.503	-0.421	0.447	-0.056	-0.014	0.024	0.011
X11	0.175	0.097	0.041	-0.707	0.100	0.668	-0.034	-0.025	0.011	0.011	-0.005
Eigenvalue	3.366	2.270	1.726	1.147	0.775	0.651	0.627	0.343	0.050	0.029	0.016
Proportion	30.60	20.64	15.700	10.42	7.05	5.92	5.70	3.12	0.45	0.27	0.15
Cumulative (%)	30.60	51.23	66.93	77.35	84.40	90.31	96.01	99.13	99.58	99.85	100

(0.43**), number of grains per spike (0.33**), biomass (0.82**), plant height (0.41**), HI (0.48**), canopy temperature depression (0.20*) and membrane thermo stability (0.18*) whereas, negative significant correlation with days to heading (-0.23**) and days to maturity (-0.23**) (Table 1). Number of tillers per plant showed significant correlations with number of grains per plant (0.78**), biomass (0.65**) and plant height (0.25**), while negative significant correlations with grain weight per spike (-0.40) and grains per spike (-0.29**). Number of grains per plant appeared to an important trait as it showed significant positive associations

with all the traits except days to heading and maturity where it was significantly negatively correlated. Grain weight per spike and number of grains per spike were also related to majority of the traits except canopy temperature depression and membrane thermo stability indicating their limited role in heat stress related parameters. Biomass per plant showed significantly correlated with plant height (0.41**) and canopy temperature depression. Plant height showed negative correlations with days to heading (-0.35**) and days to maturity (-0.33**). Harvest index was positively related with the grain yield and majority of its components but non significantly

Table 6: Normal Root-Mean-Square Distance of cluster formation

Number of cluster	Clusters Joined		Frequency	Normal Root-Mean- Square Distance
11	X3	X7	2	0.0107
10	X1	X10	2	0.0174
9	Y	CL10	3	0.0258
8	CL9	CL11	5	0.0529
7	X6	X8	2	0.1200
6	CL7	X11	3	0.1649
5	CL8	X5	6	0.2073
4	CL6	X9	4	0.3066
3	CL5	X4	7	0.3943
2	CL3	CL4	11	0.6766
1	CL2	X2	12	2.1672

Table 7: Path coefficient analysis of grain yield per plant with its component characters in recombinant inbred lines of bread wheat

Variables	Effects via											Direct effect	Indirect effect	Genotypic correlation coefficients
	X1	X2	X3	X4	X5	X6	X7	X8	X9	X10	X11			
X1	0.292	0.299	-0.278	0.155	0.153	0.017	0.000	-0.010	0.008	0.001	0.001	0.292	0.346	0.638**
X2	0.160	0.544	0.155	-0.306	0.140	0.019	0.057	-0.028	0.023	0.001	0.002	0.544	0.223	0.767**
X3	-0.125	0.130	0.647	-0.413	0.023	0.011	0.080	-0.032	0.026	<0.001	<0.001	0.647	-0.2996	0.3474**
X4	-0.084	0.310	0.498	-0.537	0.019	0.006	0.070	-0.027	0.022	0.000	<0.001	-0.537	0.8152	0.2782**
X5	0.210	0.359	0.071	-0.048	0.213	0.031	-0.003	-0.028	0.021	0.001	0.001	0.213	0.615	0.828**
X6	0.075	0.163	0.116	-0.053	0.100	0.066	0.003	-0.048	0.036	<0.001	0.0001	0.066	0.3919	0.4579**
X7	0.000	0.184	0.311	-0.225	-0.004	0.001	0.167	-0.026	0.019	<0.001	0.001	0.167	0.2613	0.4283**
X8	-0.023	-0.119	-0.161	0.113	-0.044	-0.024	-0.033	0.130	-0.103	<0.001	-0.002	0.130	-0.3956	-0.2656**
X9	-0.023	-0.119	-0.161	0.113	-0.042	-0.022	-0.030	0.126	-0.106	<0.001	-0.002	-0.106	-0.1598	-0.2658**
X10	0.052	0.141	0.038	-0.080	0.044	-0.003	0.013	0.013	-0.006	0.004	0.002	0.004	0.214	0.218**
X11	0.052	0.114	0.012	-0.032	0.034	0.007	0.018	-0.024	0.022	0.001	0.010	0.010	0.204	0.214**

Residual Effect = 0.2030; *, **: Significant at 5% and 1% level of probability, respectively

related with canopy temperature depression and membrane thermo stability. Among the heat stress related parameters, membrane thermo stability seemed to be a reliable parameter as it showed significant negative correlations with days to heading (-0.18*), days to maturity (-0.22*) and significant positive correlation with canopy temperature depression (0.22*).

Multiple linear regression analysis

Table 2 demonstrated regression coefficients and the probability of the estimated variables in predicting wheat genotypes grain yield under terminal heat stress conditions. In view of following equation.

$$Y = -4.53647 + 0.00537X2 + 1.21177X3 - 0.03024X4 + 0.21503X5 + 21.89501X7.$$

The equation explained almost 98% of the total variation within the grain yield components and the remaining 2 % assigned to residual effects. The t-test showed that number of tillers per plant (X1), numbers of grains per plant (X2), grain weight per spike (X3), number of grains per spike (X4), biomass (X5) and HI (X7) significantly contributed the grain yield. These results showed the importance of the mentioned variables in breeding programs of wheat under terminal heat stress conditions.

Stepwise multiple linear regression analysis

Data presented in Table 3 shows the partial and cumulative R² and the probability for the accepted limiting three variables in grain yield prediction. These variables are biomass (69.63%), harvest index (27.91 %), days to maturity (0.05%). According to the results 97.59% of the total variation in grain yield could be attributed to these aforementioned three variables.

Regression coefficients for the accepted variables are shown in Table 4. The equation for prediction of grain yield was obtained as given below:

$$Y\text{-intercept (a)} = "6.22311 + 0.28057X5 + 27.70354X7 - 0.01382X9$$

Principal component analysis

Results of the principal component analysis (Table 3 and Fig. 2) revealed that an increase in the number of components was associated with a decrease in eigen values. This trend reached its maximum at five factors. Accordingly, the principal components analysis had grouped the estimated wheat variables into six main components which all together accounted for 84.4% of the total variation of grain yield under terminal heat stress conditions. The first principal component which accounted for 30.6% of total variation in grain yield and mainly contributed by the characters numbers of grains/plant (0.430), numbers of grain/spike (0.323) and biomass (0.351). The second principal component made 20.64% contribution towards total variation in grain yield and represented mainly by the characters, namely, numbers of tillers/plant (0.576) and biomass (0.392). The contribution of third principal component was 15.70% and represented mainly by numbers of grains/plant (0.300), days to heading (0.525), days to maturity (0.512) and canopy temperature depression (0.404). The fourth principal component accounted 10.42% towards total variation and mainly responsible for plant height (0.390). The contribution of fifth principal component was 7.05 and it was mainly responsible for harvest index (0.721).

Cluster analysis

(Dhanda and Munjal, 2012). Positive associations of biomass (Riaz-Ud-Din *et al.* 2010) and HI (Sud and Bhagwat, 2010; Riaz-Ud-Din *et al.* 2010) with grain yield under heat stress environment indicative of efficient utilization of photosynthates. The significant correlations of grain yield with number of tillers per plant, number of grains per plant, grain weight per spike, number of grains per spike under heat stress conditions suggested contribution of component traits under heat stress (Ferrante *et al.* 2013; Slafera *et al.* 2014). The positive association of plant height with grain yield revealed that the taller plants, in general, more biomass. The results are in agreement with Jat *et al.* (2015). Days to heading and days to maturity indicated that the genotypes earlier in heading and maturity were better for grain yield and its components. These may be due to the escape from heat stress by accelerating their life cycle. These results are in line with those reported by Mondal *et al.* (2013); Zarei *et al.* (2013). Canopy temperature depression played an important role because of its positive correlations with numbers of grains per plant, numbers of grains per plant and biomass (Bahar *et al.* 2011). Membrane thermo stability also appeared to very important trait as the genotypes having high membrane thermo stability were also had high grain yield, number of grains per plant, high canopy temperature depression and earlier in days to heading and maturity. Ram *et al.* (2014) also found positive association between membrane thermo stability and grain yield. High temperature during day time particularly day temperatures above threshold level may have deleterious effects on plant physiological process (Trivedi, 2015). Therefore, the characters, namely membrane thermo stability, canopy temperature depression and earlier days to heading may help the plant to face heat stress (Satbhai *et al.* 2014).

Correlation coefficient analysis revealed that the genotypes having higher grain yield under heat stress environment were not only having higher number of yield component traits but also high membrane thermo stability, high canopy temperature depression and earlier in days to heading and maturity. Membrane thermo stability may be measured through the estimation of electrolyte leakage from the leaves of the plant when exposed to high temperature. Thus genotypes with low leakage of electrolyte may contribute heat tolerance. Canopy temperature depression also helps the plant to protect itself from heat stress by making its canopy cooler during high evaporative demand (Waiker and Arun 2015). In addition the high grain yield also contributed by the traits like earlier heading and earlier maturity under present study. Identification of suitable recombinant lines with high temperature tolerance may play an important role in improvement of wheat productivity particularly under heat stressed areas. Furthermore, the multiple statistical procedures which have been used in this study showed that simple correlation and path analysis cannot distinguish crucial variables affecting grain yield, the conclusive assertion cannot be done on the basis of these methods as such. Using multiple statistical analysis of wheat yield under stressed conditions, Leilah and Khateeb (2005) showed that weight of grains/spike, harvest index and biomass were the crucial yield attributing parameters. It seems that use multivariable statistical methods like, stepwise regression together with cluster analysis and

principal component analyses are stronger statistical methods to be used in breeding programs for screening important traits.

REFERENCES

- Al-joubri, H. A., Miller, P. A. and Robinson, H. F. 1958.** Genotype and environment variance in an upland cotton of interspecific origin. *Agron J.* **50:** 663-667.
- Bahar, B., Yildirim, Y. and Yucel, C. 2011.** Heat and drought resistance criteria in spring bread wheat (*Triticum aestivum* L.): morpho-physiological parameters for heat tolerance. *Scientific Research Essays.* **6:** 2212-2220.
- Blum, A. 1988.** Heat tolerance, In Plant Breeding for Stress Environments. *CRC Press. Inc., Boca Raton, Florida*
- Blum, A., Klueva, N. and Nguyen, H. T. 2001.** Wheat cellular thermo-tolerance is related to yield under stress. *Euphytica.* **117:** 117-123.
- Del Moral, L. G., Rharrabti, Y., Villegas, D. and Royo, C. 2003.** Evaluation of grain yield and its components in durum wheat under Mediterranean conditions. *Agron J.* **95:** 266-274.
- Dewey, D. R. and Lu, K. H. 1959.** A correlation and path coefficient analysis of crested wheat grass seed production. *Agron J.* **51:** 515-518.
- Dhanda, S. S. and Munjal, R. 2006.** Inheritance of cellular thermo-tolerance in bread wheat. *Plant Breed.* **125:** 557-567.
- Dhanda, S. S. and Munjal, R. 2012.** Heat tolerance in relation to acquired thermo-tolerance for membrane lipids in bread wheat. Association Analysis of Genomic Loci Important for Grain Weight Control in Elite Common Wheat Varieties Cultivated with Variable Water and Fertiliser Supply. *Field Crops Res.* **135:** 30-37
- Draper, N. and Smith, H. 1981.** Applied Regression Analysis, 2d Edition, New York: J. Wiley and Sons, Inc
- Everitt, B. S. 1993.** Cluster analysis. Wiley, New York. New York
- Everitt, B. S. and Dunn, G. 1992.** Applied multivariate data analysis. *Oxford University Press, New York.*
- Ferrante, A., Savin, R. and Slafer, G. A. 2013.** Floret development and grain setting differences between modern durum wheats under contrasting nitrogen availability. *Exp Bot.* **64:** 169-184.
- Gibson, L. R. and Paulsen, G. M. 1999.** Yield components of wheat grown under high temperature stress during reproductive growth. *Crop Sci.* **39:** 1841-1846.
- Ibrahim, A. M. H. and Quick, J. S. 2001.** Heritability of heat tolerance in winter and spring wheat. *Crop Sci.* **41:** 1401-1405.
- Jat, B. S., Ranwah, B. R., Baudh, Bharti and Parihar, A. K. 2015.** Stability analysis to ascertain the performance of different genotypes of wheat [*Triticum aestivum* L.] *The Bioscan.* **10(2):** 929-933.
- Kalyar, T., Rauf, S., Silva, JATD. and Shahzad, T. 2013.** Handling sunflower (*Helianthus annuus* L.) populations under heat stress. *Arch Agron. and Soil Sci.* **60:** 655-672.
- Leilah, A. A. and Al-Khateeb, S. A. 2005.** Statistical analysis of wheat yield under drought conditions. *J. Arid. Environ.* **61:** 483-496.
- Lobell, D. B., Sibley, A. and Ortiz-Monasterio, J. I. 2012.** Extreme heat effects on wheat senescence in India, *Nature. Climate Change.* **2:** 186-189.
- Modarresi, M., Mohammadi, V., Zali, A. and Mardi, M. 2010.** Response of wheat yield and yield related traits to high temperature. *Cereal Res. Commun.* **38:** 23-31.
- Mondal, S., Singh, R. P., Crossa, J., Huerta-Espino, J., Sharma, I., Chatrath, R., Singh, G. P., Sohu, V. S., Mavi, G. S., Sukuru, V. S. P., Kalappanavar, I. K., Mishra, V. K., Hussain, M., Gautam, N. R.,**

- Uddin, J., Barmam, N. C. D., Hakim, A. and Joshi, A. K . 2013.** Earliness in wheat: a key to adaptation under terminal and continual high temperature stress in South Asia. *Field Crops Res.* **151**:19-26
- Pandey, G. C., Mamrutha, H. M., Tiwari, R., Sareen, S., Bhatia, S., Siwach, P., Tiwari, V. and Sharma, I. 2015.** Physiological traits associated with heat tolerance in bread wheat (*Triticumaestivum* L.). *Physiol Mol. Biol. Plants.* **21**: 93-9.
- R Core Team 2013.** R. A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org/>.
- Ram, M., Singh, R. M. and Agrawal, R. K. 2014.** Genetic analysis for terminal heat stress in bread wheat (*Triticum aestivum* L. em thell). *The Bioscan.* **9(2)**: 771-776.
- Riaz-Ud-Din., Subhani, G. M., Ahmad, N., Hussain, M. and Rehman, A. 2010.** Effect of temperature on development and grain formation in spring wheat. *Pak. J. Bot.* **42**: 899-906.
- Saed-Moucheshi, A., Fasihfar, E., Hasheminasab, H., Rahmani, A. and Ahmadi, A. 2013.** A review on applied multivariate statistical techniques in agriculture and plant science. *Int. J. Agron. Plant. Prod.* **4**: 127-141.
- SAS Institute. 2009.** The SAS system for Windows. Release 9.2. SAS Institute, Inc., Cary, North Carolina.
- Satbhai, R., Kale, A. and Naik, R. 2014.** Cell viability, time laps study and membrane stability index during temperature induction response in wheat. *The Eco. Scan.* **VI**: 245-252.
- Savchenko, G. E., Klyuchareva, E. A., Abrabchik, L. M. and Serdyuchenko, E. V . 2002.** Effect of periodic heat shock on the membrane system of etioplasts. *Russ. J. Plant Physiol.* **49**: 349-359.
- Shiferaw, B., Smale, B., Braun, H. J., Duveiller, E., Reynolds, M. and Muricho, G. 2013 .** Crops that feed the world 10. Past successes and future challenges to the role played by wheat in global food security. *Food Security.* **5**: 291-317.
- Slafera, A. G., Savinb, R. and Sadras, V. O .2014.** Coarse and fine regulation of wheat yield components in response to genotype and environment. *Field Crops Res.* **157**: 71-83.
- Snedecor, G. W., Cochran, W. G. 1981.** Statistical methods, seventh ed. Iowa State University Press, Iowa, USA.
- Steel RGD and Torrie, J. H. 1960.** Principles and procedures of statistics, Mc Graw Hill, New York.
- Sud Suman and Bhagwat, S. G. 2010.** Assessment of acquired thermotolerance in Indian bread wheat and association with yield and component traits under heat stress environment. *Food Agri. & Envo.* **8**: 622-627.
- Sullivan, C. Y. 1972.** Mechanisms of heat and drought resistance in grain sorghum and methods of measurement. In: Rao, N. G. P., House, L.R. (Eds.), Sorghum in seventies. *Oxford & IBH Publishing Co.* New Delhi
- Trivedi, A. K. 2015.** Adaptations and Mechanisms of Heat Stress Tolerance of Plants *Academic Res. J. Agric. Sci. Res.* **3(7)**: 151-160.
- Waiker Smita and Arun, B. 2015.** Genetic diversity studies among different morpho-physiological traits in spring wheat (*Triticum aestivum* L.) in relation to heat tolerance. *The Bioscan.* **10(2)**: 911-915.
- Zarei, B., Naderi, A., Kamali, J., Lack, M. R. and Modhej, A. 2013.** Determination of physiological traits related to terminal drought and heat stress tolerance in spring wheat genotypes. *Int. J. Agric. Crop. Sci.* **5**: 2511-2520.
- Zhang, K., Wang, J., Zhang, L., Rong, C., Zhao, F., Peng, T., Li H, Cheng, D., M Liu, X., Qin, H., Zhang, A., Tong, Y. and Wang, D. 2013.** Association analysis of genomic loci important for grain weight control in elite common wheat varieties cultivated with variable water and fertiliser supply. *PLoS One.* **8**: e57853.
- Zhang, L. Y., Liu, D. C., Guo, X. L., Yang, W. L., Sun, J. Z., Wang D.W. and Zhang, A. M. 2010.** Genomic distribution of quantitative trait loci for yield and yield-related traits in common wheat. *J Integr Plant Biol.* **52**: 996-1007.
- Abbreviations:** MTS = Membrane thermo stability, CTD = Canopy temperature depression, DTM = Days to maturity, DTH = Days to heading, HI = Harvest index

